

SPECTRAL SHIFT FUNCTION OF THE SCHRÖDINGER OPERATOR  
IN THE LARGE COUPLING CONSTANT LIMIT, II.  
POSITIVE PERTURBATIONS

A. Pushnitski

Department of Mathematical Sciences  
Loughborough University  
Loughborough LE11 3TU, U.K.  
E-mail: a.b.pushnitski@lboro.ac.uk

M. Ruzhansky

Department of Mathematics  
Imperial College  
London SW7 2BZ, U.K.  
E-mail: ruzh@ic.ac.uk

**Abstract**

For the operator  $-\Delta$  in  $L^2(\mathbb{R}^d)$ ,  $d \geq 1$ , a perturbation potential  $V(x) \geq 0$ , and a coupling constant  $\alpha > 0$ , the spectral shift function  $\xi(\lambda; -\Delta + \alpha V, -\Delta)$  is considered. Assuming that  $V(x)$  decays as  $|x|^{-l}$ ,  $l > d$ , for large  $x$ , we prove that the leading term of the asymptotics of the spectral shift function as  $\alpha \rightarrow \infty$  has the form  $C\alpha^{d/l}$ , where the coefficient  $C$  is explicitly computed. The asymptotics is in agreement with the phase space volume considerations. A generalization of this result is obtained by replacing  $-\Delta$  by  $h(-\Delta)$  for a class of functions  $h$ .

## 1 INTRODUCTION

**1. Spectral Shift Function.** Let  $H$  and  $H_0$  be self-adjoint operators in a Hilbert space. Suppose that

$$H - H_0 \in \mathfrak{S}_1, \quad (1.1)$$

where  $\mathfrak{S}_1$  is the trace class. Then there exists a function  $\xi \in L^1(\mathbb{R})$  such that for all  $\phi \in C_0^\infty(\mathbb{R})$  the following *Lifshits-Krein trace formula* holds [13, 12]:

$$\mathrm{Tr}(\phi(H) - \phi(H_0)) = \int_{-\infty}^{\infty} \xi(\lambda) \phi'(\lambda) d\lambda, \quad \phi \in C_0^\infty(\mathbb{R}). \quad (1.2)$$

The function  $\xi(\lambda) = \xi(\lambda; H, H_0)$  is called the *spectral shift function* (SSF) for the pair  $H_0, H$ . The SSF is related to the scattering matrix  $S(\lambda)$  of the pair  $H_0, H$  (see, e.g., [25] for the definition of the scattering matrix) by the *Birman-Krein formula* [4]:

$$\det S(\lambda) = e^{-2\pi i \xi(\lambda)}, \quad (1.3)$$

which is valid for a.e.  $\lambda$  in the absolutely continuous spectrum of  $H_0$ . Due to (1.3), the SSF is often called *the scattering phase*. Formula (1.3) is sometimes interpreted as the definition of  $\xi$ . For an exposition of the SSF theory and the history of the subject, see [8] or [25, §8].

**2. Schrödinger Operator.** Let  $H_0 = -\Delta$  in the Hilbert space  $L^2(\mathbb{R}^d)$ ,  $d \geq 1$ , and let  $H$  be the Schrödinger operator,  $H = H_0 + V$ , where  $V$  is the operator of multiplication by a real valued potential function  $V(x)$ . It is well known (see [26] or [19, Theorem XI.12]) that under the assumption

$$|V(x)| \leq \frac{C}{(1 + |x|)^l}, \quad l > d, \quad (1.4)$$

for sufficiently large  $a > 0$  and  $k > 0$  one has

$$(H + a)^{-k} - (H_0 + a)^{-k} \in \mathfrak{S}_1. \quad (1.5)$$

This inclusion allows one to define the SSF  $\xi(\lambda; (H + a)^{-k}, (H_0 + a)^{-k})$ . The SSF for the pair  $H_0, H$  is then defined via the ‘invariance principle’:

$$\xi(\lambda; H, H_0) := -\xi((\lambda + a)^{-k}; (H + a)^{-k}, (H_0 + a)^{-k}). \quad (1.6)$$

Thus defined, the SSF  $\xi(\lambda; H, H_0)$  still obeys the trace formula (1.2). The definition (1.6) does not depend on the choice of  $k$  and  $a$ . See [25, §8.9] and [18] for an analysis of the definition (1.6) from the operator theoretic point of view. For  $\lambda < 0$ , the SSF  $\xi(\lambda; H, H_0)$  reduces to the eigenvalue counting function for the Schrödinger operator  $H$  (see, e.g., [25, §8.2]).

Various results about the high energy and semiclassical asymptotic behaviour of the SSF of the Schrödinger operator are known; see, e.g., the survey [21] and references therein. In this paper we address the less studied problem of the behaviour of the SSF in the large coupling constant limit. Namely, let  $\alpha > 0$  be a parameter (coupling constant); consider  $\xi(\lambda; H_0 + \alpha V, H_0)$  as  $\alpha \rightarrow \infty$ . For the case of non-positive perturbations  $V \leq 0$ , satisfying (1.4) with  $l > \max\{d, 2\}$ , the following formula has been proven in [17]:

$$\begin{aligned} \xi(\lambda; H_0 + \alpha V, H_0) &= -C_{1.7} \alpha^{d/2} (1 + o(1)), \quad \alpha \rightarrow \infty, \quad V \leq 0, \quad \text{a.e. } \lambda \in \mathbb{R}, \\ C_{1.7} &= (2\pi)^{-d} \omega_d \int_{\mathbb{R}^d} |V(x)|^{d/2} dx, \quad \omega_d = \text{vol}\{x \in \mathbb{R}^d \mid |x| < 1\} \end{aligned} \quad (1.7)$$

(here and elsewhere a constant which first appears in formula (i.j) is denoted by  $C_{i.j}$ ). For  $\lambda < 0$ , formula (1.7) reduces to the well-known Weyl law for the eigenvalue counting function of  $H_0 + \alpha V$  (see, e.g., [20, Theorem XIII.79]).

**3. Main Result.** The purpose of this paper is to give an asymptotic formula for the SSF  $\xi(\lambda; H_0 + \alpha V, H_0)$  for the case of non-negative perturbations  $V \geq 0$ . Here we are forced to consider potentials  $V$  with power asymptotics at infinity. More precisely, we assume that for a function  $\Psi \in C(\mathbb{S}^{d-1})$ , one has

$$\sup_{\hat{x} \in \mathbb{S}^{d-1}} |V(x) - \Psi(\hat{x})| |x|^{-l} = o(|x|^{-l}), \quad |x| \rightarrow \infty, \quad \hat{x} = \frac{x}{|x|}, \quad (1.8)$$

where  $l > d$ . In the one-dimensional case,  $\mathbb{S}^0 = \{-1, 1\}$  and  $\Psi$  reduces to a pair of numbers,  $\Psi(-1)$ ,  $\Psi(1)$ . Note that (1.8) implies (1.4). Note also that the conditions  $V \geq 0$  and (1.8) imply that  $\Psi \geq 0$ .

Our main result is

**Theorem 1.1.** *Let  $H_0 = -\Delta$  in  $L^2(\mathbb{R}^d)$ ,  $d \geq 1$  and let  $V \geq 0$  be a bounded function which satisfies (1.8) with some  $\Psi \in C(\mathbb{S}^{d-1})$ ,  $\Psi \geq 0$ , and  $l > d$ . Then, for all  $\lambda > 0$ , one has*

$$\begin{aligned} \xi(\lambda; H_0 + \alpha V, H_0) &= C_{1.9} \alpha^{d/l} (1 + o(1)), \quad \alpha \rightarrow \infty, \quad V \geq 0, \quad \lambda > 0, \\ C_{1.9} &= (2\pi)^{-d} d^{-1} \int_{|p|^2 < \lambda} (\lambda - |p|^2)^{-d/l} dp \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}. \end{aligned} \quad (1.9)$$

In the one-dimensional case,  $\int_{\mathbb{S}^0} \Psi^{d/l}(\hat{x}) d\hat{x} \equiv \Psi^{d/l}(-1) + \Psi^{d/l}(1)$ .

**4. Remarks.** 1. Computing the integral over  $p$  in (1.9), one obtains

$$C_{1.9} = \frac{1}{d} (4\pi)^{-d/2} \frac{\Gamma(1 - \frac{d}{l})}{\Gamma(1 + \frac{d}{2} - \frac{d}{l})} \lambda^{\frac{d}{2} - \frac{d}{l}} \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}.$$

2. The trace formula (1.2) defines  $\xi(\lambda)$  as an element of  $L^1(\mathbb{R})$ . However, under the hypothesis of Theorem 1.1, the spectral shift function  $\xi(\lambda)$  is actually

continuous in  $\lambda > 0$ . This can be seen e.g., by employing (1.3) and using the fact that the scattering matrix is a continuous function of the spectral parameter in the trace class topology.

Under more restrictive conditions on  $V$ , one can obtain results on smoothness of the SSF (see, e.g. [21] and references therein).

3. Formula (1.9) explains why for  $V \geq 0$  we are forced to consider the more restrictive class (1.8) of potentials  $V$  (as compared to (1.4)): the order  $d/l$  of  $\alpha$  in (1.9) is determined by the asymptotic behaviour of  $V$ . Note that the order  $d/2$  of  $\alpha$  in (1.7) does not depend on  $V$ . In a certain sense, the order of the asymptotics (1.7) is determined by the region of large momenta, whereas the order of (1.9) is determined by the region of large coordinates.

4. Following the proof, one can check that the asymptotics (1.9) is uniform in  $\lambda$  on every compact subset of  $(0, \infty)$ .

5. In the case of the potential  $V$  of a variable sign, one would expect a superposition of the terms of the type  $-C_{1.7}\alpha^{d/2}$  and  $C_{1.9}\alpha^{d/l}$  in the asymptotics of the SSF. If

$$\frac{d}{2} > \frac{d}{l}, \quad (1.10)$$

then it is natural to expect the leading term of the asymptotics to have the order  $d/2$  and be ‘generated’ by the negative part of  $V$ . This is indeed the case; it has been proven in [17] that for  $V$  of a variable sign, under the assumption (1.10) the asymptotic formula (1.7) holds (in the sense of  $L^1_{\text{loc}}$  convergence) with  $C_{1.7}$  replaced by

$$(2\pi)^{-d}\omega_d \int_{V(x)<0} |V(x)|^{d/2} dx.$$

More precise (in terms of the type of convergence) results can be obtained by using the technique of [22].

Note that, since  $l > d$ , the inequality (1.10) always holds if  $d \geq 2$ .

6. In [24], the following bound for the SSF has been obtained:

$$|\xi(\lambda; H_0 + \alpha V, H_0)| \leq C\lambda^{\frac{d}{2} - \frac{d}{l}} \alpha^{d/l}, \quad V \geq 0. \quad (1.11)$$

Theorem 1.1 shows that this bound has a sharp order in  $\alpha$  and  $\lambda$ . Note that the proof of (1.11) in [24] was based on a hypothesis (Assumption 4.3 in [24]) which has not been proven so far. In fact, the main technical result of the present paper (Theorem 4.4) provides the proof of this hypothesis.

7. The asymptotic coefficient  $C_{1.9}$  can be interpreted in terms of the phase space volume. An elementary calculation shows that

$$C_{1.9} = (2\pi)^{-d} \lim_{\alpha \rightarrow \infty} \alpha^{-d/l} \text{vol}\{(x, p) \in \mathbb{R}^{2d} \mid |p|^2 < \lambda < |p|^2 + \alpha V(x)\}.$$

Another compact form for  $C_{1.9}$  is

$$C_{1.9} = \int_{\mathbb{R}^d} (\tau(\lambda) - \tau(\lambda - \Psi(\hat{x})|x|^{-l})) dx, \quad (1.12)$$

where  $\tau$  is the integrated density of states of  $-\Delta$ ,

$$\tau(\lambda) = (2\pi)^{-d} \text{vol}\{p \in \mathbb{R}^d \mid |p|^2 < \lambda\} = (2\pi)^{-d} \omega_d (\max\{\lambda, 0\})^{d/2}.$$

Without going into details, let us mention that the asymptotics (1.9) with the coefficient  $C_{1.9}$  expressed in the form (1.12) formally coincides with the asymptotics of the eigenvalue counting function in the gaps of a perturbed periodic Schrödinger operator [2, Theorem 2.1].

**5. Generalization.** In Theorem 1.1, one can replace the operator  $H_0 = -\Delta$  by the operator  $H_0 = h(-\Delta)$ , where  $h$  is a sufficiently regular function. More precisely, we assume that

$$h \in C^2(\mathbb{R}), \quad h(0) = 0, \quad h'(r) > 0 \quad \forall r > 0, \quad (1.13)$$

$$h(r) \geq Cr^\gamma, \quad C > 0, \quad \gamma > 0 \quad \text{for all large } r > 0. \quad (1.14)$$

Define  $H_0 = h(-\Delta)$  according to the spectral theorem, and let  $H = H_0 + V$ , where  $V$  obeys (1.4). By (1.14), the relation (1.5) holds for large enough  $a > 0$  and  $k > 0$  (see [26]). Thus, the SSF  $\xi(\lambda; H, H_0)$  is well defined via the ‘invariance principle’ (1.6).

**Theorem 1.2.** *Let  $h$  satisfy (1.13) and (1.14), and let  $H_0 = h(-\Delta)$  in  $L^2(\mathbb{R}^d)$ ,  $d \geq 1$ . Let  $V \geq 0$  be a bounded function which satisfies (1.8) with some  $\Psi \in C(\mathbb{S}^{d-1})$ ,  $\Psi \geq 0$ , and  $l > d$ . Then, for all  $\lambda > 0$ , one has*

$$\begin{aligned} \xi(\lambda; H_0 + \alpha V, H_0) &= C_{1.15} \alpha^{d/l} (1 + o(1)), \quad \alpha \rightarrow \infty, \\ C_{1.15} &= (2\pi)^{-d} d^{-1} \int_{h(|p|^2) < \lambda} (\lambda - h(|p|^2))^{-d/l} dp \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}. \end{aligned} \quad (1.15)$$

The constant  $C_{1.15}$  can be written in the form (1.12) with

$$\tau(\lambda) = (2\pi)^{-d} \text{vol}\{p \in \mathbb{R}^d \mid h(|p|^2) < \lambda\}.$$

In order to prove Theorem 1.2, one needs to make only minor modifications in the proof of Theorem 1.1. In Sections 2–8 we explicitly prove Theorem 1.1 and in the end of Section 7 we comment on these modifications. One can relax the assumptions (1.13), (1.14) in various directions.

**6. Notation.** We denote  $D = -i\nabla$ ; in this notation,  $H_0 = D^2$ . We will use the notation  $\chi(x^2 < a)$ ,  $\chi(x^2 > a)$ , etc. for the characteristic function of  $\{x \in \mathbb{R}^d \mid |x|^2 < a\}$ ,  $\{x \in \mathbb{R}^d \mid |x|^2 > a\}$ , etc. Then,  $\chi(D^2 < a)$ ,  $\chi(D^2 > a)$ ,

etc. are the spectral projections of  $D^2$  corresponding to the intervals  $(-\infty, a)$ ,  $(a, \infty)$ , etc. We write  $\langle x \rangle = (1 + |x|^2)^{1/2}$ . For  $p \in \mathbb{R}^d$ , we shall write  $p^2$  instead of  $|p|^2$  for brevity. We do not distinguish between a function of  $x \in \mathbb{R}^d$  and an operator of multiplication by this function in  $L^2(\mathbb{R}^d)$ . We assume the following convention about the signs  $\pm$ : if in any statement we use double indices ' $\pm$ ', then this statement should be interpreted as a pair of statements, in one of which all the indices take the value '+', and in another one — the value '-'. For a real-valued function  $\mu$ , we denote  $\mu_{\pm} = (|\mu| \pm \mu)/2$ . Finally, by  $C$  (possibly with subscripts) we denote constants, whose value is of no importance for us; these constants may depend on  $d, l, V, \lambda$  and other auxiliary parameters, and this dependence is suppressed in the notation.

7. Let us describe the technique and the structure of the paper. Our main operator theoretic tool is a formula representation from [16] for the spectral shift function. This representation reduces the proof of Theorem 1.1 to the problem of computing the asymptotics of the spectrum of the sandwiched resolvent  $\sqrt{V}(H_0 - \lambda - i0)^{-1}\sqrt{V}$ . This reduction is carried out in Section 2. In Section 3 for the ease of further reference we collect the necessary information about the weak Schatten classes of compact operators.

In the rest of the paper, Sections 4–8, we compute the asymptotics of the spectrum of the operator  $\sqrt{V}(H_0 - \lambda - i0)^{-1}\sqrt{V}$ . The main difficulty here is the singularity of the symbol of  $(H_0 - \lambda - i0)^{-1}$ . In Section 4 we consider the operator  $\sqrt{V}(H_0 - \lambda)^{-1}\chi(|H_0 - \lambda| > \varepsilon)\sqrt{V}$  ( $\varepsilon > 0$  small) with the singularity 'removed'; for this operator the required asymptotics can be obtained by employing the result of [6]. In Sections 5–8, we show that removing the singularity as above does not affect the spectral asymptotics. In this part of the paper, we find it convenient to use the language of variational quotients (VQ); the translation of the problem into this language is given in Section 5.

## 2 REPRESENTATION FOR THE SSF

Let  $H_0$  and  $V$  be as in Theorem 1.1. Here and in the rest of the paper we use the notation  $W = \sqrt{V}$ . For  $\text{Im } z > 0$  introduce the compact operators

$$T(z) = W(H_0 - z)^{-1}W,$$

$$A(z) = \text{Re } T(z) = (T(z) + T(z)^*)/2, \quad B(z) = \text{Im } T(z) = (T(z) - T(z)^*)/(2i).$$

By (1.4), *the limit  $T(\lambda + i0)$  exists in the operator norm for all  $\lambda > 0$*  (see [1] or [20, §XIII.8]); here one needs  $l > 1$  rather than  $l > d$ . The operators  $A(\lambda) \equiv A(\lambda + i0)$  and  $B(\lambda) \equiv B(\lambda + i0)$  are compact and self-adjoint.

For any compact self-adjoint operator  $M$ , we use the notation

$$n_{\pm}(s, M) = \#\{n \mid \pm\lambda_n(M) > s\}, \quad s > 0, \quad (2.1)$$

where  $\lambda_n(M)$  are the eigenvalues of  $M$ , enumerated with the multiplicities taken into account. Similarly,

$$n(s, M) = n_+(s, M) + n_-(s, M) = \#\{n \mid |\lambda_n(M)| > s\}, \quad s > 0. \quad (2.2)$$

The following representation for the SSF is valid:

$$\xi(\lambda; H_0 + \alpha V, H_0) = \frac{1}{\pi} \int_{-\infty}^{\infty} n_-(\alpha^{-1}, A(\lambda) + tB(\lambda))(1 + t^2)^{-1} dt. \quad (2.3)$$

The representation (2.3) has been found in [16] in the framework of general operator theory. Application to the Schrödinger operator has been discussed in [17]. A similar formula is valid for negative perturbations:

$$\xi(\lambda; H_0 - \alpha V, H_0) = -\frac{1}{\pi} \int_{-\infty}^{\infty} n_+(\alpha^{-1}, A(\lambda) + tB(\lambda))(1 + t^2)^{-1} dt.$$

A more general formula representation is available now (see [10, 18]), which is applicable to perturbations  $V$  of a variable sign.

Formula (2.3) allows one to reduce the problem of computing the asymptotics of  $\xi(\lambda; H_0 + \alpha V, H_0)$  as  $\alpha \rightarrow \infty$  to computing the asymptotics of the spectra of compact self-adjoint operators  $A(\lambda)$ ,  $B(\lambda)$ . This follows from the simple lemma below, which has been proven in [17].

**Lemma 2.1.** [17, Lemma 4.4] *Let  $A$  and  $B$  be compact self-adjoint operators such that for some  $\varkappa > \sigma > 0$ ,  $\sigma < 1$ , one has*

$$\lim_{s \rightarrow 0} s^{\varkappa} n_-(s, A) = C_{2.4}, \quad \limsup_{s \rightarrow 0} s^{\sigma} n(s, B) < \infty. \quad (2.4)$$

Then

$$\lim_{\alpha \rightarrow \infty} \alpha^{-\varkappa} \frac{1}{\pi} \int_{-\infty}^{\infty} n_-(\alpha^{-1}, A + tB)(1 + t^2)^{-1} dt = C_{2.4}.$$

The rest of the present paper is devoted to the proof of the following two relations:

$$\lim_{s \rightarrow 0} s^{d/l} n_-(s, A(\lambda)) = C_{1.9}, \quad (2.5)$$

$$\lim_{s \rightarrow 0} s^{\frac{d-1}{l-1}} n(s, B(\lambda)) < \infty, \quad d \geq 2. \quad (2.6)$$

From here Theorem 1.1 will follow. Indeed, for  $d \geq 2$  one has  $\frac{d-1}{l-1} < \frac{d}{l}$ , and thus Lemma 2.1 (with  $\varkappa = d/l$ ,  $\sigma = (d-1)/(l-1)$ ) together with (2.5), (2.6) gives the proof of Theorem 1.1. For  $d = 1$  it is easy to see that  $\text{rank } B(\lambda) = 2$ , and therefore

$$\limsup_{s \rightarrow 0} s^{\sigma} n(s, B(\lambda)) < \infty \quad \text{for any } \sigma > 0, \quad d = 1.$$

Thus Theorem 1.1 will follow from Lemma 2.1 (with  $\varkappa = 1/l$  and any  $\sigma \in (0, 1/l)$ ) and (2.5).

### 3 CLASSES $\Sigma_{\varkappa}$ OF COMPACT OPERATORS

In this section we collect the necessary information about the weak Schatten classes  $\Sigma_{\varkappa}$  of compact operators. We use the paper [5] as a main source of reference; however, most of the material mentioned here (apart from Proposition 3.3 and the inequalities (3.5), (3.6)) can be found, for example, in the monograph [11] (see also [23]).

**1.** For a compact operator  $A$  in a Hilbert space, its singular numbers are denoted by  $s_n(A) = \sqrt{\lambda_n(A^*A)}$ . For  $\varkappa > 0$ , the class  $\Sigma_{\varkappa}$  consists of all compact operators  $A$  such that

$$|A|_{\varkappa} \equiv \sup_n n^{1/\varkappa} s_n(A) < \infty. \quad (3.1)$$

The functional (3.1) defines a quasinorm in  $\Sigma_{\varkappa}$ . With respect to this quasinorm, the space  $\Sigma_{\varkappa}$  is non-separable (unless the underlying Hilbert space is finite-dimensional) and complete. For  $\varkappa > 1$  the space  $\Sigma_{\varkappa}$  can be equipped with a norm, which is different from, but equivalent to the quasinorm  $|\cdot|_{\varkappa}$ ; we will not need the explicit formula for this norm. From the definition of  $s_n(A)$  it follows that

$$A^*A \in \Sigma_{\varkappa/2} \quad \Leftrightarrow \quad A \in \Sigma_{\varkappa}. \quad (3.2)$$

**2.** Similarly to (2.1) and (2.2), for a compact (but not necessarily self-adjoint) operator  $A$  one denotes

$$n(s, A) = \#\{n \mid s_n(A) > s\}, \quad s > 0.$$

Note that this notation is consistent with (2.2), as for a compact self-adjoint operator  $A$  one has  $s_n(A) = |\lambda_n(A)|$ .

For a compact operator  $A$  one can define the following functionals:

$$\begin{aligned} \Delta_{\varkappa}(A) &= \limsup_{s \rightarrow 0} s^{\varkappa} n(s, A), \\ \delta_{\varkappa}(A) &= \liminf_{s \rightarrow 0} s^{\varkappa} n(s, A). \end{aligned} \quad (3.3)$$

For  $A \in \Sigma_{\varkappa}$  one has

$$s^{\varkappa} n(s, A) \leq |A|_{\varkappa}^{\varkappa}, \quad s > 0,$$

and so the functionals (3.3) are finite on  $\Sigma_{\varkappa}$ . Conversely, if  $\Delta_{\varkappa}(A) < \infty$ , then  $A \in \Sigma_{\varkappa}$ .

For a compact self-adjoint operator  $A$ , one can define also the following functionals:

$$\begin{aligned} \Delta_{\varkappa}^{(\pm)}(A) &= \limsup_{s \rightarrow 0} s^{\varkappa} n_{\pm}(s, A), \\ \delta_{\varkappa}^{(\pm)}(A) &= \liminf_{s \rightarrow 0} s^{\varkappa} n_{\pm}(s, A), \end{aligned} \quad (3.4)$$

where  $n_{\pm}$  is defined in (2.1). The functionals (3.4) are finite on  $\Sigma_{\varkappa}$ . If  $\Delta_{\varkappa}^{(+)}(A) < \infty$  and  $\Delta_{\varkappa}^{(-)}(A) < \infty$ , then  $A \in \Sigma_{\varkappa}$ . However, any one of the conditions  $\Delta_{\varkappa}^{(+)}(A) < \infty$ ,  $\Delta_{\varkappa}^{(-)}(A) < \infty$  alone does not imply that  $A \in \Sigma_{\varkappa}$ ; in fact, later we will deal with the case  $\Delta_{\varkappa}^{(-)}(A) < \infty$ ,  $A \notin \Sigma_{\varkappa}$  — see Remark 4.6 below.

**3.** The functionals  $\Delta_{\varkappa}^{(\pm)}$ ,  $\delta_{\varkappa}^{(\pm)}$  are continuous in  $\Sigma_{\varkappa}$ . This follows from the inequalities (see [5], eq. (4), (6))

$$\begin{aligned} |(\Delta_{\varkappa}^{(\pm)}(A_2))^{1/(\varkappa+1)} - (\Delta_{\varkappa}^{(\pm)}(A_1))^{1/(\varkappa+1)}| \\ \leq (\Delta_{\varkappa}(A_2 - A_1))^{1/(\varkappa+1)} \leq |A_2 - A_1|_{\varkappa}^{\varkappa/(\varkappa+1)}, \end{aligned} \quad (3.5)$$

$$\begin{aligned} |(\delta_{\varkappa}^{(\pm)}(A_2))^{1/(\varkappa+1)} - (\delta_{\varkappa}^{(\pm)}(A_1))^{1/(\varkappa+1)}| \\ \leq (\Delta_{\varkappa}(A_2 - A_1))^{1/(\varkappa+1)} \leq |A_2 - A_1|_{\varkappa}^{\varkappa/(\varkappa+1)}. \end{aligned} \quad (3.6)$$

Similarly, the functionals  $\Delta_{\varkappa}$  and  $\delta_{\varkappa}$  are continuous in  $\Sigma_{\varkappa}$ .

If  $A$  is compact and  $B_1, B_2$  are bounded, then  $s_n(B_1AB_2) \leq \|B_1\| \|B_2\| s_n(A)$ . It follows that

$$A \in \Sigma_{\varkappa}, \quad B_1, B_2 \text{ are bounded} \Rightarrow B_1AB_2 \in \Sigma_{\varkappa}, \quad (3.7)$$

$$|B_1AB_2|_{\varkappa} \leq \|B_1\| \|B_2\| |A|_{\varkappa}. \quad (3.8)$$

Similarly, if  $A = A^*$  is compact and  $B$  is bounded, then

$$\Delta_{\varkappa}^{(\pm)}(B^*AB) \leq \|B\|^{2\varkappa} \Delta_{\varkappa}^{(\pm)}(A), \quad (3.9)$$

$$\delta_{\varkappa}^{(\pm)}(B^*AB) \leq \|B\|^{2\varkappa} \delta_{\varkappa}^{(\pm)}(A). \quad (3.10)$$

Write  $A_1 \geq A_2$  if  $(A_1f, f) \geq (A_2f, f)$  for all vectors  $f$ . Due to the variational properties of eigenvalues, the functionals  $\Delta_{\varkappa}^{(\pm)}$ ,  $\delta_{\varkappa}^{(\pm)}$  are monotone with respect to the relation ‘ $\geq$ ’. In particular, we will need the following statement:

$$A_1 \geq A_2 \quad \Rightarrow \quad \Delta_{\varkappa}^{(-)}(A_1) \leq \Delta_{\varkappa}^{(-)}(A_2). \quad (3.11)$$

**4.** The class  $\Sigma_{\varkappa}^0 \subset \Sigma_{\varkappa}$  consists of all operators  $A$  such that  $s_n(A) = o(n^{-1/\varkappa})$ . One can prove that  $\Sigma_{\varkappa}^0$  is closed in  $\Sigma_{\varkappa}$ . The space  $\Sigma_{\varkappa}^0$  is separable and coincides with the closure in  $\Sigma_{\varkappa}$  of the set of all finite rank operators. It is easy to see that  $A \in \Sigma_{\varkappa}^0$  if and only if  $\Delta_{\varkappa}(A) = 0$ . One has

$$A^*A \in \Sigma_{\varkappa/2}^0 \quad \Leftrightarrow \quad A \in \Sigma_{\varkappa}^0. \quad (3.12)$$

If  $A \in \Sigma_{\nu}$ , then  $A \in \Sigma_{\varkappa}^0$  for any  $\varkappa > \nu$ .

The A. Horn’s inequality for singular numbers of a product of two operators (see [14] or [11]) yields

**Proposition 3.1.** *If  $A_1 \in \Sigma_{\varkappa}$  and  $A_2 \in \Sigma_{\nu}$ , then  $A_1A_2 \in \Sigma_{\mu}$ , where  $\frac{1}{\mu} = \frac{1}{\varkappa} + \frac{1}{\nu}$ . If, moreover,  $A_1 \in \Sigma_{\varkappa}^0$  or  $A_2 \in \Sigma_{\nu}^0$ , then  $A_1A_2 \in \Sigma_{\mu}^0$ .*

The following statement (which first appeared in [7]) is a modification of the well known result (due to H. Weyl and K. Fan) about the asymptotics of the singular numbers of a sum of two operators (see, e.g., [11], Theorem 2.3).

**Proposition 3.2.** *Let  $A_1, A_2$  be self-adjoint operators of the class  $\Sigma_{\varkappa}$ . If  $A_1 - A_2 \in \Sigma_{\varkappa}^0$ , then*

$$\Delta_{\varkappa}^{(\pm)}(A_1) = \Delta_{\varkappa}^{(\pm)}(A_2), \quad \delta_{\varkappa}^{(\pm)}(A_1) = \delta_{\varkappa}^{(\pm)}(A_2).$$

The proof can be obtained by a direct application of (3.5), (3.6).

5. Finally, we will need the following statement from [5].

**Proposition 3.3.** *[5, Theorem 3] Let  $A_1, A_2$  be self-adjoint operators of the class  $\Sigma_{\varkappa}$ . Suppose that  $\Delta_{\varkappa}^{(\pm)}(A_1) = \delta_{\varkappa}^{(\pm)}(A_1)$ ,  $\Delta_{\varkappa}^{(\pm)}(A_2) = \delta_{\varkappa}^{(\pm)}(A_2)$ , and  $A_1^*A_2 \in \Sigma_{\varkappa/2}^0$ ,  $A_1A_2^* \in \Sigma_{\varkappa/2}^0$ . Then*

$$\Delta_{\varkappa}^{(\pm)}(A_1 + A_2) = \delta_{\varkappa}^{(\pm)}(A_1 + A_2) = \Delta_{\varkappa}^{(\pm)}(A_1) + \Delta_{\varkappa}^{(\pm)}(A_2).$$

#### 4 PROOF OF (2.5) AND (2.6)

In this section we reduce the proof of (2.5) and (2.6) to several simpler statements, which we will prove in the rest of the paper.

First, we state a result which is a ‘spherically symmetric version’ of (2.6).

**Lemma 4.1.** *Let  $d \geq 2$ . For any  $l > d$  and  $\lambda > 0$ , one has*

$$\operatorname{Im} (\langle x \rangle^{-l/2} (D^2 - \lambda - i0)^{-1} \langle x \rangle^{-l/2}) \in \Sigma_{\frac{d-1}{l-1}}. \quad (4.1)$$

We prove Lemma 4.1 in Sections 5, 6.

**Proof of (2.6):** Write  $W(x) = X(x) \langle x \rangle^{-l/2}$  with  $X$  bounded. Now (2.6) follows from (3.7) and Lemma 4.1. ■

Next, we state a result from [6] (Theorem 2 and following this theorem Remarks 1–3).

**Proposition 4.2.** *Let  $a, b \in L^\infty(\mathbb{R}^d)$ ,  $a, b$  have compact support. Let  $V \geq 0$  be a bounded function which satisfies (1.8) with some  $\Psi \in C(\mathbb{S}^{d-1})$ ,  $\Psi \geq 0$ , and  $l > 0$ . Then*

$$a(D)Vb(D) \in \Sigma_{d/l}, \quad (4.2)$$

$$\begin{aligned} \Delta_{d/l}(a(D)Vb(D)) &= \delta_{d/l}(a(D)Vb(D)) \\ &= (2\pi)^{-d} d^{-1} \int_{\mathbb{R}^d} (a(p)b(p))^{d/l} dp \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}. \end{aligned} \quad (4.3)$$

In fact, (4.2) and (4.3) have been proven in [6] for a much broader class of  $a$  and  $b$ , but the above statement is sufficient for our purposes.

Note that, taking  $a(D) = b(D)$  in (4.2) and using (3.2), we get

$$a(D)W \in \Sigma_{2d/l}. \quad (4.4)$$

We will need a slightly different version of Proposition 4.2:

**Proposition 4.3.** *Let  $\mu \in L^1(\mathbb{R}^d)$ ,  $\mu$  have a compact support. Let  $V$  be as in Theorem 1.1 and  $W = \sqrt{V}$ . Then  $W\mu(D)W \in \Sigma_{d/l}$  and*

$$\begin{aligned} \Delta_{d/l}^{(\pm)}(W\mu(D)W) &= \delta_{d/l}^{(\pm)}(W\mu(D)W) \\ &= (2\pi)^{-d}d^{-1} \int_{\mathbb{R}^d} (\mu_{\pm}(p))^{d/l} dp \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}. \end{aligned} \quad (4.5)$$

*Proof.* 1. First assume that  $\mu \geq 0$ . Then

$$\lambda_n(W\mu(D)W) = \lambda_n(\mu^{1/2}(D)V\mu^{1/2}(D)),$$

and the result follows from Proposition 4.2.

2. For  $\mu$  of a variable sign, write

$$W\mu(D)W = W\mu_+(D)W - W\mu_-(D)W.$$

By the previous step and Proposition 3.3, it suffices to check that

$$(W\mu_+(D)W)(W\mu_-(D)W) \in \Sigma_{d/(2l)}^0. \quad (4.6)$$

3. Write the operator in (4.6) as

$$(W\mu_+^{1/2}(D))(\mu_+^{1/2}(D)V\mu_-^{1/2}(D))(\mu_-^{1/2}(D)W).$$

By (4.4), the first and the third term in the above product belong to  $\Sigma_{2d/l}$ . By (4.3), the second term belongs to  $\Sigma_{d/l}^0$  (as  $\mu_+\mu_- \equiv 0$  and the integral in (4.3) vanishes). Thus, applying Proposition 3.1 to the above product, we get (4.6). ■

Next, we state the main technical result of the paper.

**Theorem 4.4.** *For any  $l > d$  and  $\lambda > 0$ , one has*

$$\operatorname{Re}(\langle x \rangle^{-l/2}(D^2 - \lambda - i0)^{-1}\chi(D^2 < 2\lambda)\langle x \rangle^{-l/2}) \in \Sigma_{d/l}, \quad (4.7)$$

$$\Delta_{d/l}(\operatorname{Re}(\langle x \rangle^{-l/2}(D^2 - \lambda - i0)^{-1}\chi(|D^2 - \lambda| < \varepsilon)\langle x \rangle^{-l/2})) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (4.8)$$

We prove this theorem in Sections 5, 7–8. This theorem will allow us to get rid of the singularity of the symbol of  $A(\lambda)$ . Note that (4.7) has been conjectured in [24, Assumption 4.3].

Finally, we will need the following technical result.

**Proposition 4.5.** *Let  $V$  be as in Theorem 1.1 and  $W = \sqrt{V}$ . Then, for any  $\lambda > 0$ , one has*

$$\chi(D^2 < \lambda)W\chi(D^2 > \lambda) \in \Sigma_{2d/l}^0. \quad (4.9)$$

We postpone the proof of this proposition till the end of this section.

**Proof of (2.5):** 1. We will prove separately the following two inequalities:

$$\Delta_{d/l}^{(-)}(A(\lambda)) \leq C_{1.9}, \quad (4.10)$$

$$\delta_{d/l}^{(-)}(A(\lambda)) \geq C_{1.9}. \quad (4.11)$$

Clearly, (4.10) and (4.11) imply (2.5).

2. In order to prove (4.10), we first notice that

$$A(\lambda) \geq \operatorname{Re}(W(D^2 - \lambda - i0)^{-1}\chi(D^2 < 2\lambda)W).$$

By (3.11), this yields

$$\Delta_{d/l}^{(-)}(A(\lambda)) \leq \Delta_{d/l}^{(-)}(\operatorname{Re}(W(D^2 - \lambda - i0)^{-1}\chi(D^2 < 2\lambda)W)).$$

Next, we factorize  $W(x) = \langle x \rangle^{-l/2}X(x)$  with bounded  $X$  and for any  $\varepsilon > 0$  write

$$\begin{aligned} & \operatorname{Re}(W(D^2 - \lambda - i0)^{-1}\chi(D^2 < 2\lambda)W) \\ &= W(D^2 - \lambda)^{-1}\chi(D^2 < 2\lambda)\chi(|D^2 - \lambda| > \varepsilon)W \\ &+ X\operatorname{Re}(\langle x \rangle^{-l/2}(D^2 - \lambda - i0)^{-1}\chi(|D^2 - \lambda| < \varepsilon)\langle x \rangle^{-l/2})X. \end{aligned}$$

By (4.8) and (3.9),

$$\Delta_{d/l}(X\operatorname{Re}(\langle x \rangle^{-l/2}(D^2 - \lambda - i0)^{-1}\chi(|D^2 - \lambda| < \varepsilon)\langle x \rangle^{-l/2})X) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

By (3.5), it follows

$$\Delta_{d/l}^{(-)}(A(\lambda)) = \Delta_{d/l}^{(-)}(W(D^2 - \lambda)^{-1}\chi(D^2 < 2\lambda)\chi(|D^2 - \lambda| > \varepsilon)W) + o(1), \quad \varepsilon \rightarrow 0.$$

Finally, by Proposition 4.3,

$$\begin{aligned} & \Delta_{d/l}^{(-)}(W(D^2 - \lambda)^{-1}\chi(D^2 < 2\lambda)\chi(|D^2 - \lambda| > \varepsilon)W) \\ &= (2\pi)^{-d}d^{-1} \int_{p^2 < \lambda - \varepsilon} (\lambda - p^2)^{-d/l} dp \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$ , we get (4.10).

3. Let us prove (4.11). Denote for brevity  $E_1 = \chi(D^2 < 2\lambda)$ ,  $E_2 = \chi(D^2 > 2\lambda)$ . By (3.10),

$$\delta_{d/l}^{(-)}(A(\lambda)) \geq \delta_{d/l}^{(-)}(E_1 A(\lambda) E_1). \quad (4.12)$$

For any  $\varepsilon > 0$ , write

$$\begin{aligned} E_1 A(\lambda) E_1 &= E_1 W (D^2 - \lambda)^{-1} \chi(|D^2 - \lambda| > \varepsilon) W E_1 \\ &\quad + E_1 \operatorname{Re}(W (D^2 - \lambda - i0)^{-1} \chi(|D^2 - \lambda| < \varepsilon) W) E_1. \end{aligned} \quad (4.13)$$

By (4.4) with  $a(D) = \chi(D^2 < 2\lambda)$ , one has  $E_1 W \in \Sigma_{2d/l}$ . Thus, by Proposition 3.1, the first term in the r.h.s. of (4.13) belongs to  $\Sigma_{d/l}$ . The second term belongs to  $\Sigma_{d/l}$  by Theorem 4.4. Thus,  $E_1 A(\lambda) E_1 \in \Sigma_{d/l}$ . Next, as on the previous step of the proof, using (4.8), (3.6) and (3.9), we get

$$\delta_{d/l}^{(-)}(E_1 A(\lambda) E_1) = \delta_{d/l}^{(-)}(E_1 W (D^2 - \lambda)^{-1} \chi(|D^2 - \lambda| > \varepsilon) W E_1) + o(1), \quad \varepsilon \rightarrow 0. \quad (4.14)$$

4. Denote  $\mu(D) = (D^2 - \lambda)^{-1} \chi(|D^2 - \lambda| > \varepsilon)$ . Let us check that

$$E_1 W \mu(D) W E_1 - W E_1 \mu(D) E_1 W \in \Sigma_{d/l}^0. \quad (4.15)$$

One has

$$\begin{aligned} E_1 W \mu(D) W E_1 - W E_1 \mu(D) E_1 W &= (E_1 W E_2) \mu(D) (E_2 W E_1) \\ &\quad - (E_2 W E_1) \mu(D) (E_1 W E_1) - (E_1 W E_1) \mu(D) (E_1 W E_2) - (E_2 W E_1) \mu(D) (E_1 W E_2). \end{aligned}$$

By Proposition 4.5 and Proposition 3.1, all the terms in the r.h.s. of the last equation belong to  $\Sigma_{d/l}^0$ .

5. By (4.15), Proposition 3.2 and Proposition 4.3,

$$\begin{aligned} \delta_{d/l}^{(-)}(E_1 W (D^2 - \lambda)^{-1} \chi(|D^2 - \lambda| > \varepsilon) W E_1) \\ &= \delta_{d/l}^{(-)}(W E_1 (D^2 - \lambda)^{-1} \chi(|D^2 - \lambda| > \varepsilon) E_1 W) \\ &= (2\pi)^{-d} d^{-1} \int_{p^2 < \lambda - \varepsilon} (\lambda - p^2)^{-d/l} dp \int_{\mathbb{S}^{d-1}} \Psi^{d/l}(\hat{x}) d\hat{x}. \end{aligned}$$

This, together with (4.12) and (4.14), gives (4.11). ■

**Remark 4.6.** Note that (see [17])  $A(\lambda) \in \Sigma_{d/2}$  and  $\Delta_{d/2}^{(+)}(A(\lambda)) \neq 0$ . Thus,  $A(\lambda)$  does *not* belong to  $\Sigma_{d/l}$ , unless  $d/l \geq d/2$  (this can happen only for  $d = 1$ ).

*Proof of Proposition 4.5:* 1. First, as above, denote for brevity  $E_1 = \chi(D^2 < 2\lambda)$ ,  $E_2 = \chi(D^2 > 2\lambda)$ . Next, fix a function  $\zeta$ ,

$$\begin{aligned} \zeta &\in C^\infty(\mathbb{R}^d), \quad \zeta(x) \geq 0 \text{ for all } x \in \mathbb{R}^d, \\ \zeta(x) &= 1 \text{ for } |x| \geq 2, \quad \zeta(x) = 0 \text{ for } |x| \leq 1. \end{aligned} \quad (4.16)$$

Denote  $W_0(x) = \Psi^{1/2}(\widehat{x})|x|^{-l/2}\zeta(x)$ . By (1.8), for any  $\varepsilon > 0$  one can write  $W = W_0 + W_\varepsilon + \widetilde{W}_\varepsilon$ , where  $\widetilde{W}_\varepsilon$  has a compact support and

$$|W_\varepsilon(x)| \leq \varepsilon|x|^{-l/2}\zeta(x). \quad (4.17)$$

According to this decomposition, write

$$E_1 W E_2 = E_1 W_0 E_2 + E_1 W_\varepsilon E_2 + E_1 \widetilde{W}_\varepsilon E_2. \quad (4.18)$$

First note that by Proposition 4.2,  $E_1 \widetilde{W}_\varepsilon^2 E_1 \in \Sigma_{d/l}^0$ ; thus,  $E_1 \widetilde{W}_\varepsilon \in \Sigma_{2d/l}^0$  and so the first term in the r.h.s. of (4.18) belongs to  $\Sigma_{2d/l}^0$ . Next, by (4.4) and (4.17),

$$|E_1 W_\varepsilon E_2|_{2d/l} \leq |E_1 W_\varepsilon|_{2d/l} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Thus, since  $\Sigma_{2d/l}^0$  is closed in  $\Sigma_{2d/l}$ , it is sufficient to prove that

$$E_1 W_0 E_2 \in \Sigma_{2d/l}^0. \quad (4.19)$$

2. By (3.8),

$$|E_1 W_0 E_2|_{2d/l} \leq |E_1|x|^{-l/2}\zeta(x)|_{2d/l} \|\Psi^{1/2}\|_{L^\infty(\mathbb{S}^{d-1})}.$$

Thus, approximating  $\Psi^{1/2}$  in  $C(\mathbb{S}^{d-1})$  by smooth functions, we approximate the operator (4.19) in  $\Sigma_{2d/l}$ . Since  $\Sigma_{2d/l}^0$  is closed in  $\Sigma_{2d/l}$ , it is sufficient to prove (4.19) for smooth  $\Psi^{1/2}$  (in fact, we will need  $\Psi^{1/2} \in C^1(\mathbb{S}^{d-1})$ ).

3. By Proposition 4.2,

$$\chi(D^2 < \lambda)W_0\chi(\lambda < D^2 < \lambda + R) \in \Sigma_{2d/l}^0,$$

as  $\chi(p^2 < \lambda)\chi(\lambda < p^2 < \lambda + R) \equiv 0$  and the integral in (4.3) vanishes. Thus, it suffices to prove that

$$\chi(D^2 < \lambda)W_0\chi(D^2 > \lambda + R) \in \Sigma_{2d/l}^0 \quad (4.20)$$

for all large enough  $R > 0$ .

4. The operator in (4.20) is unitarily equivalent to the operator

$$\chi(|x| < \sqrt{\lambda})W_0(D)\chi(|x| > \sqrt{\lambda + R}). \quad (4.21)$$

Take  $R \geq 4 + 4\sqrt{\lambda}$ , so that  $\sqrt{\lambda + R} \geq \sqrt{\lambda} + 2$ . Then the integral kernel  $K(x, y)$  of the operator (4.21) vanishes for  $|x - y| \leq 2$ . Thus, multiplying the kernel  $K(x, y)$  by  $\zeta(x - y)$  (where  $\zeta$  is as in (4.16)) does not change the operator (4.21).

The last statement can be rephrased as follows. Let  $\eta(x) = 1 - \zeta(x)$ , let  $\widehat{\eta}$  be the (unitary) Fourier transform of  $\eta$ , and let  $W_0 * \widehat{\eta}$  be the convolution of  $W_0$  and  $\widehat{\eta}$ . Then

$$\begin{aligned} & \chi(|x| < \sqrt{\lambda})W_0(D)\chi(|x| > \sqrt{\lambda} + 2) \\ &= \chi(|x| < \sqrt{\lambda})(W_0(D) - (2\pi)^{-d/2}(W_0 * \widehat{\eta})(D))\chi(|x| > \sqrt{\lambda} + 2) \end{aligned} \quad (4.22)$$

Below we prove that

$$\lim_{|x| \rightarrow \infty} |x|^{l/2} |W_0(x) - (2\pi)^{-d/2}(W_0 * \widehat{\eta})(x)| = 0. \quad (4.23)$$

By Proposition 4.2, this implies

$$\chi(|x| < \sqrt{\lambda})(W_0(D) - (2\pi)^{-d/2}(W_0 * \widehat{\eta})(D)) \in \Sigma_{2d/l}^0.$$

By (4.22), this yields (4.20).

5. To complete the proof, it remains to check (4.23). This is a straightforward computation. First note that  $\widehat{\eta}$  is in the Schwartz class. Next,  $\eta(0) = 1$  and so

$$(2\pi)^{-d/2} \int_{\mathbb{R}^d} \widehat{\eta}(y) dy = 1.$$

Thus,

$$\begin{aligned} & |x|^{l/2} (W_0(x) - (2\pi)^{-d/2}(W_0 * \widehat{\eta})(x)) \\ &= (2\pi)^{-d/2} |x|^{l/2} \int_{\mathbb{R}^d} \widehat{\eta}(y) (W_0(x) - W_0(x-y)) dy \\ &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \widehat{\eta}(y) (\Psi^{1/2}(\widehat{x})\zeta(x) - \Psi^{1/2}(\widehat{x-y})|x|^{l/2}|x-y|^{-l/2}\zeta(x-y)) dy. \end{aligned}$$

Let us split the last integral into the sum of two integrals over  $|y| \leq \sqrt{|x|}$  and  $|y| \geq \sqrt{|x|}$  and check that both integrals tend to zero as  $|x| \rightarrow \infty$ . For the first integral, one has

$$\begin{aligned} & \left| \int_{|y| \leq \sqrt{|x|}} (\dots) dy \right| \leq \|\widehat{\eta}\|_{L^1} \|\Psi^{1/2}\|_{L^\infty} \sup_{|y| \leq \sqrt{|x|}} | |x|^{l/2} |x-y|^{-l/2} \zeta(x-y) - \zeta(x) | \\ & \quad + \|\widehat{\eta}\|_{L^1} \sup_{|y| \leq \sqrt{|x|}} | \Psi^{1/2}(\widehat{x-y}) - \Psi^{1/2}(\widehat{x}) | = O(1/\sqrt{|x|}), \quad |x| \rightarrow +\infty. \end{aligned}$$

In the last estimate we have used the fact that  $\Psi^{1/2} \in C^1(\mathbb{S}^{d-1})$ .

Next,

$$\begin{aligned} & \left| \int_{|y| \geq \sqrt{|x|}} (\dots) dy \right| \leq \left( \int_{|y| \geq \sqrt{|x|}} |\widehat{\eta}(y)| dy \right) \\ & \quad \times \sup_{|y| \geq \sqrt{|x|}} | \Psi^{1/2}(\widehat{x-y}) |x|^{l/2} |x-y|^{-l/2} \zeta(x-y) - \Psi^{1/2}(\widehat{x})\zeta(x) |; \end{aligned}$$

here the first term decays faster than any power of  $|x|$ , and the second term is  $O(|x|^{l/2})$ . ■

## 5 VARIATIONAL QUOTIENTS

In the rest of the paper, we prove Lemma 4.1 and Theorem 4.4. In this section, we translate Lemma 4.1 and Theorem 4.4 into the language of variational quotients (VQ).

Let  $\mu \in L^1(\mathbb{R}^d)$  be a real-valued compactly supported function. Consider the VQ

$$t[u] = \frac{\int_{\mathbb{R}^d} \mu(p) |u(p)|^2 dp}{\|u\|_{H^{l/2}}^2}, \quad u \in \text{Dom}(t) = H^{l/2}(\mathbb{R}^d), \quad (5.1)$$

where  $l > d$ . Here the norm in the Sobolev space  $H^{l/2}(\mathbb{R}^d)$  is defined by

$$\|u\|_{H^{l/2}}^2 = \int_{\mathbb{R}^d} \langle p \rangle^l |\widehat{u}(p)|^2 dp, \quad (5.2)$$

where  $\widehat{u}$  is the Fourier transform of  $u$ .

Consider the numerator of (5.1) as a quadratic form in the Hilbert space  $\text{Dom}(t)$  with the metric given by the denominator of (5.1); denote this quadratic form by  $\mathfrak{t}[u, u]$ . The form  $\mathfrak{t}[u, u]$  generates a self-adjoint operator  $T$  (in the same Hilbert space):

$$\mathfrak{t}[u, u] = (Tu, u)_{H^{l/2}}, \quad u \in \text{Dom}(T),$$

where  $(\cdot, \cdot)_{H^{l/2}}$  is the inner product, corresponding to the norm (5.2):

$$(u, v)_{H^{l/2}} = \int_{\mathbb{R}^d} \langle p \rangle^l \widehat{u}(p) \overline{\widehat{v}(p)} dp. \quad (5.3)$$

The *spectrum* of the VQ (5.1) is the spectrum of the operator  $T$ . Below we use the notation of the type

$$n(s, (5.1)) := n(s, T), \quad s > 0,$$

where the r.h.s. is defined by (2.2). By the variational characterization of eigenvalues, one has

$$n(s, (5.1)) = \sup\{\dim \mathcal{L} \mid \mathcal{L} \subset \text{Dom}(t), |t[u]| > s \quad \forall u \in \mathcal{L}\};$$

here  $\mathcal{L}$  is a linear subspace with the stated properties.

It is easy to see that the operator  $T$ , generated by (5.1), is unitarily equivalent to the operator  $\langle x \rangle^{-l/2} \mu(D) \langle x \rangle^{-l/2}$  in the Hilbert space  $L^2(\mathbb{R}^d)$ . Thus,

$$n(s, \langle x \rangle^{-l/2} \mu(D) \langle x \rangle^{-l/2}) = n(s, (5.1)), \quad s > 0.$$

Fix  $\lambda > 0$  and  $\varepsilon > 0$ . Taking  $\mu(p) = \chi(|p^2 - \lambda| < \varepsilon) \operatorname{Re}(p^2 - \lambda - i\delta)^{-1}$  and letting  $\delta \rightarrow +0$ , we see that the VQ

$$\frac{v.p. \int_{|p^2 - \lambda| < \varepsilon} (p^2 - \lambda)^{-1} |u(p)|^2 dp}{\|u\|_{H^{l/2}}^2}, \quad u \in H^{l/2}(\mathbb{R}^d) \quad (5.4)$$

determines the spectrum of the operator from (4.8):

$$n(s, \operatorname{Re}(\langle x \rangle^{-l/2} (D^2 - \lambda - i0)^{-1} \chi(|D^2 - \lambda| < \varepsilon) \langle x \rangle^{-l/2})) = n(s, (5.4)). \quad (5.5)$$

In the same way, the VQ

$$\frac{(\pi/2) \lambda^{(d/2)-1} \int_{\mathbb{S}^{d-1}} |u(\sqrt{\lambda}\omega)|^2 d\omega}{\|u\|_{H^{l/2}}^2}, \quad u \in H^{l/2}(\mathbb{R}^d) \quad (5.6)$$

determines the spectrum of the operator from (4.1):

$$n(s, \operatorname{Im}(\langle x \rangle^{-l/2} (D^2 - \lambda - i0)^{-1} \langle x \rangle^{-l/2})) = n(s, (5.6)). \quad (5.7)$$

In Section 6, we prove that

$$\limsup_{s \rightarrow 0} s^{\frac{d-1}{l-1}} n(s, (5.6)) < \infty. \quad (5.8)$$

By (5.7), this yields Lemma 4.1.

In Sections 7–8 we prove that

$$\limsup_{s \rightarrow 0} s^{d/l} n(s, (5.4)) \leq C_{5.9} = (2\pi)^{-d} \omega_d \int_{|p^2 - \lambda| < \varepsilon} |p^2 - \lambda|^{-d/l} dp. \quad (5.9)$$

By (5.5), this yields Theorem 4.4.

In dealing with variational quotients, we will use the following simple facts:

(i) Increasing the VQ leads to the increase of the counting function  $n$ :

$$t_1[u] \leq t_2[u] \quad \forall u \in \operatorname{Dom}(t_1) = \operatorname{Dom}(t_2) \quad \Rightarrow \quad n(s, T_1) \leq n(s, T_2) \quad \forall s > 0.$$

(ii) Extending the domain of the VQ leads to the increase of the counting function  $n$ :

$$t_1[u] = t_2[u] \quad \forall u \in \operatorname{Dom}(t_1) \subset \operatorname{Dom}(t_2) \quad \Rightarrow \quad n(s, T_1) \leq n(s, T_2) \quad \forall s > 0.$$

## 6 PROOF OF (5.8)

Throughout this section,  $d \geq 2$ . The key idea of the proof of (5.8) is to reduce the VQ (5.6) to a VQ on a sphere in  $\mathbb{R}^d$ .

1. First, we need notation for some sets in  $\mathbb{R}^d$ ; fix  $\lambda > 0$  and denote

$$\begin{aligned} S_\lambda &= \{x \in \mathbb{R}^d \mid |x| = \sqrt{\lambda}\}, & S_\lambda^c &= \{x \in \mathbb{R}^d \mid |x| \neq \sqrt{\lambda}\}, \\ S_\lambda^i &= \{x \in \mathbb{R}^d \mid |x| < \sqrt{\lambda}\}, & S_\lambda^e &= \{x \in \mathbb{R}^d \mid |x| > \sqrt{\lambda}\} \end{aligned} \quad (6.1)$$

(the superscripts ‘c’, ‘i’, ‘e’ stand for ‘complement’, ‘interior’, ‘exterior’). In what follows, for  $d \geq 2$  we use the Sobolev classes  $H^p(S_\lambda)$ ,  $p \in \mathbb{R}$ . The exact choice of the norm in  $H^p(S_\lambda)$  will be of no importance for us; in all the construction, this norm can be replaced by any equivalent norm. One of the possible choices of the norm is

$$\|u\|_{H^p(S_\lambda)} = \|(I - \Delta_B)^{p/2}u\|_{L^2(S_\lambda)}, \quad (6.2)$$

where  $\Delta_B$  is the Laplace-Beltrami operator on  $S_\lambda$ .

**Proposition 6.1.** *Fix  $q \in \mathbb{R}$  and  $p > 0$ ,  $p > q$ . Then, for the VQ*

$$\frac{\|u\|_{H^q(S_\lambda)}^2}{\|u\|_{H^p(S_\lambda)}^2}, \quad u \in H^p(S_\lambda), \quad (6.3)$$

one has

$$\limsup_{s \rightarrow 0} s^\varkappa n(s, (6.3)) < \infty, \quad \varkappa = \frac{d-1}{2(p-q)}. \quad (6.4)$$

*Proof:* For the choice (6.2) of the metric, the spectrum of the VQ (6.3) coincides with the spectrum of  $(I - \Delta_B)^{q-p}$ . Thus,

$$n(s, (6.3)) = n(s^{1/(p-q)}, (I - \Delta_B)^{-1}),$$

and (6.4) follows from the well-known asymptotics

$$\lim_{s \rightarrow 0} s^{(d-1)/2} n(s, (I - \Delta_B)^{-1}) = C(d). \quad \blacksquare$$

In this section, we will only use Proposition 6.1 for  $p = (l-1)/2$ ,  $q = 0$ , but we will need the general version in Section 8.

2. Denote by  $\gamma$  the following restriction operator:

$$\gamma : H^{l/2}(\mathbb{R}^d) \rightarrow H^{(l-1)/2}(S_\lambda), \quad u \mapsto u|_{S_\lambda}.$$

As it is well known, the operator  $\gamma$  is bounded:

$$\|\gamma u\|_{H^{(l-1)/2}} \leq C_{6.5} \|u\|_{H^{l/2}}. \quad (6.5)$$

This enables us to estimate the VQ (5.6) as follows:

$$\frac{\|\gamma u\|_{L^2(\mathbb{S}^{d-1})}^2}{\|u\|_{H^{l/2}(\mathbb{R}^d)}^2} \leq C_{6.5}^{-2} \frac{\|\gamma u\|_{L^2(\mathbb{S}^{d-1})}^2}{\|\gamma u\|_{H^{(l-1)/2}(\mathbb{S}^{d-1})}^2}, \quad u \in H^{l/2}(\mathbb{R}^d).$$

Thus, the spectrum of (6.3) with  $p = (l-1)/2$ ,  $q = 0$  gives the estimate for the spectrum of (5.6) and so (5.8) follows from Proposition 6.1.

## 7 PROOF OF (5.9)

1. As above, we use the notation (6.1) (for  $d \geq 1$ ) and consider  $H^{l/2}(\mathbb{R}^d)$  as a Hilbert space with respect to the inner product (5.3). Let  $H_0^{l/2}(S_\lambda^c)$  be the closure of  $C_0^\infty(S_\lambda^c)$  in  $H^{l/2}(\mathbb{R}^d)$  and let  $H_\perp^{l/2}(\mathbb{R}^d)$  be the orthogonal complement of  $H_0^{l/2}(S_\lambda^c)$ :

$$H^{l/2}(\mathbb{R}^d) = H_0^{l/2}(S_\lambda^c) \oplus H_\perp^{l/2}(\mathbb{R}^d). \quad (7.1)$$

According to (7.1), we shall write

$$u = u_0 + u_\perp, \quad u_0 \in H_0^{l/2}(S_\lambda^c), \quad u_\perp \in H_\perp^{l/2}(\mathbb{R}^d).$$

2. We will need the following two statements.

**Proposition 7.1.** *Let  $H_\perp^{l/2}(\mathbb{R}^d)$  be as defined above, and let  $\delta \geq 0$ ,  $\delta < (l-1)/2$ . Then, for the VQ*

$$\frac{\|u_\perp\|_{H^{\frac{1}{2}+\delta}(\mathbb{R}^d)}^2}{\|u_\perp\|_{H^{l/2}(\mathbb{R}^d)}^2}, \quad u_\perp \in H_\perp^{l/2}(\mathbb{R}^d), \quad (7.2)$$

one has

$$\lim_{s \rightarrow 0} s^\varkappa n(s, (7.2)) = 0, \quad \forall \varkappa > \frac{d-1}{l-2\delta-1}. \quad (7.3)$$

We prove Proposition 7.1 in Section 8.

**Proposition 7.2.** [9] *Let, as above,  $l > d$  and let  $\mu$  be a real valued compactly supported function on  $\mathbb{R}^d$  such that*

$$|\mu(p)| \leq C(|p| - \sqrt{\lambda})^{-\rho}, \quad \rho < l/d. \quad (7.4)$$

Then for any  $u_0 \in H_0^{l/2}(S_\lambda^c)$ , the integral  $\int_{\mathbb{R}^d} \mu(p) |u_0(p)|^2 dp$  converges and for the VQ

$$\frac{\int_{\mathbb{R}^d} \mu(p) |u_0(p)|^2 dp}{\|u_0\|_{H^{l/2}(\mathbb{R}^d)}^2}, \quad u_0 \in H_0^{l/2}(S_\lambda^c), \quad (7.5)$$

one has

$$\lim_{s \rightarrow 0} s^{d/l} n(s, (7.5)) = (2\pi)^{-d} \omega_d \int_{\mathbb{R}^d} |\mu(p)|^{d/l} dp. \quad (7.6)$$

For  $\mu \in L^1(\mathbb{R}^d)$ , Proposition 7.2 is well known (see, e.g, [7]). In this case (7.6) is also true (and is equivalent to Proposition 4.3 with  $V(x) = \langle x \rangle^{-l}$ ) if  $H_0^{l/2}(S_\lambda^c)$  is replaced by  $H^{l/2}(\mathbb{R}^d)$  in (7.5).

In the form, stated above, Proposition 7.2 allows for a non-integrable singularity of  $\mu$ . This situation has been studied in [3] for  $d = 1$  and in [9] for  $d \geq 2$ . Here the space  $H_0^{l/2}(S_\lambda^c)$  cannot be replaced by  $H^{l/2}(\mathbb{R}^d)$ . The construction of [3, 9] is based on the generalized Hardy inequality, which ensures that the functions  $u_0(p)$  of  $H_0^{l/2}(S_\lambda^c)$  tend to zero as  $p$  approaches  $S_\lambda$ .

Finally, note that in [9], the VQ of the type (7.5) was studied for a cube in  $\mathbb{R}^d$ , instead of the sphere  $S_\lambda$ . However, Proposition 7.2 in the above stated form follows by application of the standard technique of the local change of coordinates.

**3.** We start proving (5.9). Let us estimate the numerator of (5.4). Fix  $\tau > 0$  and  $\theta_0 > 0, \theta_1 > 0$  so that  $\theta_0 + \theta_1 = 1, \theta_0 < l/(2d), \theta_1 < 1/2$ . Then, using the decomposition (7.1) on the first step and the Cauchy inequality on the second step, we get

$$\begin{aligned}
& v.p. \int_{|p^2-\lambda|<\varepsilon} (p^2 - \lambda)^{-1} |u(p)|^2 dp = \int_{|p^2-\lambda|<\varepsilon} (p^2 - \lambda)^{-1} |u_0(p)|^2 dp \\
& + v.p. \int_{|p^2-\lambda|<\varepsilon} (p^2 - \lambda)^{-1} |u_\perp(p)|^2 dp + 2\text{Re} \int_{|p^2-\lambda|<\varepsilon} (p^2 - \lambda)^{-1} u_\perp(p) \overline{u_0(p)} dp \\
& \leq \int_{|p^2-\lambda|<\varepsilon} |p^2 - \lambda|^{-1} |u_0(p)|^2 dp + v.p. \int_{|p^2-\lambda|<\varepsilon} (p^2 - \lambda)^{-1} |u_\perp(p)|^2 dp \\
& + \tau \int_{|p^2-\lambda|<\varepsilon} |p^2 - \lambda|^{-2\theta_0} |u_0(p)|^2 dp + \frac{1}{\tau} \int_{|p^2-\lambda|<\varepsilon} |p^2 - \lambda|^{-2\theta_1} |u_\perp(p)|^2 dp. \quad (7.7)
\end{aligned}$$

As it will follow from further reasoning, all the integrals in the r.h.s., apart from the second one, converge absolutely.

Below we consider separately the four VQ, corresponding to the four terms in the r.h.s. of (7.7).

**4.** Consider the VQ

$$\frac{\int_{|p^2-\lambda|<\varepsilon} |p^2 - \lambda|^{-1} |u_0(p)|^2 dp}{\|u\|_{H^{l/2}}^2}, \quad u \in H^{l/2}(\mathbb{R}^d). \quad (7.8)$$

First note that, according to (7.1), the compact self-adjoint operator in  $H^{l/2}(\mathbb{R}^d)$ , generated by (7.8), splits into the orthogonal sum of the operator, generated by

$$\frac{\int_{|p^2-\lambda|<\varepsilon} |p^2 - \lambda|^{-1} |u_0(p)|^2 dp}{\|u_0\|_{H^{l/2}}^2}, \quad u_0 \in H_0^{l/2}(S_\lambda^c) \quad (7.9)$$

in  $H_0^{l/2}(S_\lambda^c)$  and zero operator in  $H_\perp^{l/2}(\mathbb{R}^d)$ . Thus,

$$n(s, (7.8)) = n(s, (7.9)).$$

The weight function  $\mu(p) = |p^2 - \lambda|^{-1} \chi(|p^2 - \lambda| < \varepsilon)$  in the numerator of (7.9) satisfies (7.4) and thus, by Proposition 7.2,

$$\lim_{s \rightarrow 0} s^{d/l} n(s, (7.9)) = C_{5.9}. \quad (7.10)$$

**5.** Consider the VQ, generated by the third term in (7.7). By the same reasoning as above, it reduces to the VQ in  $H_0^{l/2}(S_\lambda^c)$ :

$$\frac{\tau \int_{|p^2 - \lambda| < \varepsilon} |p^2 - \lambda|^{-2\theta_0} |u_0(p)|^2 dp}{\|u_0\|_{H^{l/2}}^2}, \quad u_0 \in H_0^{l/2}(S_\lambda^c). \quad (7.11)$$

Again, by Proposition 7.2,

$$\lim_{s \rightarrow 0} s^{d/l} n(s, (7.11)) = \tau^{d/l} (2\pi)^{-d} \omega_d \int_{|p^2 - \lambda| < \varepsilon} |p^2 - \lambda|^{-2\theta_0 d/l} dp. \quad (7.12)$$

**6.** Consider the VQ, generated by the fourth term in (7.7):

$$\frac{\tau^{-1} \int_{|p^2 - \lambda| < \varepsilon} |p^2 - \lambda|^{-2\theta_1} |u_\perp(p)|^2 dp}{\|u\|_{H^{l/2}}^2}, \quad u \in H^{l/2}(\mathbb{R}^d). \quad (7.13)$$

First note that, by the same reasoning as above, the VQ (7.13) reduces to a VQ in  $H_\perp^{l/2}(\mathbb{R}^d)$ . Next, let us estimate the numerator. For  $r > 0$ , denote

$$\tilde{u}_\perp(r) = r^{d-1} \int_{\mathbb{S}^{d-1}} |u_\perp(r\omega)|^2 d\omega. \quad (7.14)$$

Then, separating the variables in the integral, we get

$$\begin{aligned} \frac{1}{\tau} \int_{|p^2 - \lambda| < \varepsilon} |p^2 - \lambda|^{-2\theta_1} |u_\perp(p)|^2 dp &= \frac{1}{\tau} \int_{\sqrt{\lambda - \varepsilon}}^{\sqrt{\lambda + \varepsilon}} |r^2 - \lambda|^{-2\theta_1} \tilde{u}_\perp(r) dr \\ &\leq \frac{1}{\tau} \|\tilde{u}_\perp\|_{L^\infty} \int_{\sqrt{\lambda - \varepsilon}}^{\sqrt{\lambda + \varepsilon}} |r^2 - \lambda|^{-2\theta_1} dr, \end{aligned}$$

where the norm of  $\tilde{u}_\perp$  is taken in  $L^\infty(\sqrt{\lambda - \varepsilon}, \sqrt{\lambda + \varepsilon})$ . By the embedding theorem (see, e.g., [15]),

$$\|\tilde{u}_\perp\|_{L^\infty} \leq C \|u_\perp\|_{H^{1/2}(\mathbb{R}^d)}^2 \quad (7.15)$$

(here the fact that  $u_\perp$  belongs to the subspace  $H_\perp^{l/2}(\mathbb{R}^d)$  is, of course, irrelevant to the validity of (7.15); the same estimate holds for any  $u \in H^{l/2}(\mathbb{R}^d)$ ). Thus, we have arrived at the VQ (7.2) with  $\delta = 0$ ; by Proposition 7.1,

$$\lim_{s \rightarrow 0} s^\varkappa n(s, (7.13)) = 0, \quad \forall \varkappa > \frac{d-1}{l-1}. \quad (7.16)$$

**7.** Finally, consider the VQ, generated by the second integral in the r.h.s. of (7.7):

$$\frac{v.p. \int_{|p^2 - \lambda| < \varepsilon} (p^2 - \lambda)^{-1} |u_\perp(p)|^2 dp}{\|u\|_{H^{l/2}}^2}, \quad u_0 \in H^{l/2}(\mathbb{R}^d). \quad (7.17)$$

As above, it reduces to the same VQ in the space  $H_\perp^{l/2}(\mathbb{R}^d)$ . Again using the notation (7.14) and estimating the integral in the numerator of (7.17), we get

$$v.p. \int_{|p^2 - \lambda| < \varepsilon} (p^2 - \lambda)^{-1} |u_\perp(p)|^2 dp = \int_{\sqrt{\lambda - \varepsilon}}^{\sqrt{\lambda + \varepsilon}} (r^2 - \lambda)^{-1} \tilde{u}_\perp(r) dr \leq C \|\tilde{u}_\perp\|_{C^{\delta_0}}, \quad (7.18)$$

where the norm of  $\tilde{u}_\perp$  is taken in the Hölder space  $C^{\delta_0}(\sqrt{\lambda - \varepsilon}, \sqrt{\lambda + \varepsilon})$ , and the exponent  $\delta_0$  is any positive number. The following estimate holds true for all  $\delta > \delta_0$ :

$$\|\tilde{u}_\perp\|_{C^{\delta_0}} \leq C \|u_\perp\|_{H^{\frac{1}{2} + \delta}}^2. \quad (7.19)$$

This estimate is well known to specialists in function theory; however, we could not find it in textbooks on Sobolev spaces. Using the standard technique of local change of coordinates, (7.19) can be reduced to a similar estimate with a hyperplane instead of the sphere  $S_\lambda$ ; in the latter case, the proof of (7.19) is a matter of a simple calculation (similar to the one used in the proof of the embedding theorem (7.15)). The fact that  $u_\perp$  belongs to the subspace  $H_\perp^{l/2}(\mathbb{R}^d)$  is irrelevant to the validity of (7.19).

We use (7.19) and choose  $\delta$  and  $\delta_0$  in such a way that  $\frac{d-1}{l-2\delta-1} < d/l$ . Thus, we arrive at the VQ (7.2) and so by Proposition 7.1,

$$\lim_{s \rightarrow 0} s^\varkappa n(s, (7.17)) = 0, \quad \forall \varkappa > \frac{d-1}{l-2\delta-1}. \quad (7.20)$$

**8.** The relations (7.7), (7.10), (7.12), (7.16), (7.20) yield

$$\lim_{s \rightarrow 0} s^{d/l} n(s, (5.4)) \leq C_{5.9} + C\tau^{d/l};$$

since  $\tau > 0$  can be taken arbitrary small, we get (5.9). ■

**9. Remarks on the proof of Theorem 1.2.** In order to prove Theorem 1.2, one needs to make only minor modifications in the above construction. Let us briefly comment on this point. The representation (2.3) is still valid due

to the abstract result of [16, Theorem 1.2] (here the condition (1.14) is important). Next, the proof again reduces to the verification of the relations (2.5) (with  $C_{1.15}$  instead of  $C_{1.9}$ ) and (2.6). Further, in the same way the above question reduces to the consideration of variational quotients. A straightforward calculation shows that

$$\begin{aligned}\operatorname{Im}(h(p^2) - \lambda - i0)^{-1} &= \frac{1}{h'(h^{-1}(\lambda))} \operatorname{Im}(p^2 - h^{-1}(\lambda) - i0)^{-1}, \\ \operatorname{Re}(h(p^2) - \lambda - i0)^{-1} &= F_\lambda(p^2) \operatorname{Re}(p^2 - h^{-1}(\lambda) - i0)^{-1},\end{aligned}$$

where the function

$$F_\lambda(x) := \frac{x - h^{-1}(\lambda)}{h(x) - \lambda}$$

is positive,  $C^1$ -smooth and bounded away from zero in the neighbourhood of  $h^{-1}(\lambda)$ . Thus, we get extra factors in the numerators of the VQ (5.4) and (5.6).

The extra factor in (5.6) does not affect further considerations in any way. The extra factor in (5.4) results in extra factors in the integrands in the numerators of (7.8), (7.11), (7.13), and (7.17). This will only affect constants in the estimates. Most importantly, one has to use the Hölder continuity of  $F_\lambda$  when proving the analogue of (7.18).

## 8 PROOF OF PROPOSITION 7.1

The idea of the proof of Proposition 7.1 is to reduce the VQ (7.2) to a VQ on a sphere similarly to the construction of Section 6. However, here the technical details are more complicated.

**1. Change of function in (7.2).** The condition

$$u_\perp \perp H_0^{l/2}(S_\lambda^c) \quad (\text{orthogonality in } H^{l/2}) \quad (8.1)$$

can be stated as

$$((1 - \Delta)^{l/4} u_\perp, (1 - \Delta)^{l/4} \varphi)_{L^2} = 0, \quad \forall \varphi \in C_0^\infty(S_\lambda^c).$$

Write  $l/2 = m - \theta$ , where  $m \in \mathbb{Z}$ ,  $m - 1 < l/2 \leq m$ . Denote  $v = (1 - \Delta)^{-\theta} u_\perp$ ,  $v \in H^{(l/2)+2\theta}(\mathbb{R}^d)$ . Then the above condition can be stated in terms of  $v$  as

$$((1 - \Delta)^{(l/4)+\theta} v, (1 - \Delta)^{l/4} \varphi)_{L^2} = 0, \quad \forall \varphi \in C_0^\infty(S_\lambda^c),$$

which is equivalent to the differential equation ( $(l/2) + \theta = m$  is integer!)

$$(1 - \Delta)^m v(x) = 0, \quad x \in S_\lambda^c.$$

Thus, the spectrum of (7.2) coincides with the spectrum of the VQ

$$\frac{\|v\|_{H^{\frac{1}{2}+\delta+2\theta}(\mathbb{R}^d)}^2}{\|v\|_{H^{(l/2)+2\theta}(\mathbb{R}^d)}^2}, \quad v \in H^{(l/2)+2\theta}(\mathbb{R}^d), \quad (1 - \Delta)^m v(x) = 0, \quad x \in S_\lambda^c. \quad (8.2)$$

The implicit orthogonality condition (8.1) has been reduced to an elliptic differential equation on  $v$ . Obviously, if  $l$  happens to be an integer even number, the above change of function becomes vacuous: in this case,  $\theta = 0$ ,  $m = l/2$  and  $v = u_\perp$ .

For  $d = 1$ , the solutions to the above differential equation form a *finite dimensional* space, and Proposition 7.1 follows immediately. Thus, below we consider the case  $d \geq 2$ .

**2. Separation of the interior and exterior.** Here our aim is to ‘split’ the VQ (8.2) into two VQ’s: one in  $S_\lambda^i$  and another in  $S_\lambda^e$ . Below we are dealing with the Sobolev spaces  $H^s(S_\lambda^i)$  and  $H^s(S_\lambda^e)$ ,  $s > 0$ ; the exact choice of one of the possible equivalent norms in these spaces will not be important for our construction. Denote by  $P_i$  and  $P_e$  the restriction operators

$$\begin{aligned} P_i &: H^s(\mathbb{R}^d) \rightarrow H^s(S_\lambda^i), \quad u \mapsto u|_{S_\lambda^i}, \quad s > 0, \\ P_e &: H^s(\mathbb{R}^d) \rightarrow H^s(S_\lambda^e), \quad u \mapsto u|_{S_\lambda^e}, \quad s > 0. \end{aligned}$$

The operators  $P_i$  and  $P_e$  are bounded for any  $s > 0$  and thus

$$\|P_i v\|_{H^{(l/2)+2\theta}(S_\lambda^i)}^2 + \|P_e v\|_{H^{(l/2)+2\theta}(S_\lambda^e)}^2 \leq C \|v\|_{H^{(l/2)+2\theta}(\mathbb{R}^d)}^2 \quad (8.3)$$

for any  $v \in H^{(l/2)+2\theta}(\mathbb{R}^d)$ . This estimate will allow us to ‘split’ the denominator of (8.2).

In order to ‘split’ the numerator, we will also need the bound of the opposite sign:

**Proposition 8.1.** *For any  $s > 1/2$ ,  $s - (1/2) \notin \mathbb{Z}$ , there exists a constant  $C_{8.4}$  such that for any  $v \in H^s(\mathbb{R}^d)$ ,*

$$\|v\|_{H^s(\mathbb{R}^d)}^2 \leq C_{8.4} (\|P_i v\|_{H^s(S_\lambda^i)}^2 + \|P_e v\|_{H^s(S_\lambda^e)}^2). \quad (8.4)$$

This statement is probably known to specialists, but we were unable to find it in the standard monographs on Sobolev spaces, so below we give a naive do-it-yourself proof. The assumption  $s > 1/2$  is not necessary for the validity of (8.4), but sufficient for our purposes.

Obviously, for an integer  $s$ , (8.4) becomes a trivial equality with  $C_{8.4} = 1$  (for the standard choice of norm in the Sobolev spaces). Thus, if the order  $1/2 + \delta + 2\theta$  in the numerator of the VQ (8.2) is integer, we do not need to use the estimate (8.4). By using certain freedom of choice of the parameters  $m$ ,  $\theta$

and  $\delta$ , we could make  $1/2 + \delta + 2\theta$  equal to 1 when  $d$  is odd and to 0 when  $d$  is even. However, because of various restrictions on the parameters  $\delta$  and  $\theta$ , this route would only allow us to cover the case  $l > d + 1/2$ . We omit the details of this calculation.

*Proof.* 1. Let  $\pi_e : H^s(S_\lambda^e) \rightarrow H^s(\mathbb{R}^d)$  be a bounded extension operator,  $P_e \pi_e = \text{id}$ . Take  $v_e = \pi_e P_e v$ . Then

$$\|v_e\|_{H^s(\mathbb{R}^d)} \leq C \|P_e v\|_{H^s(S_\lambda^e)}. \quad (8.5)$$

2. Consider the function  $w := P_i v - P_i v_e$ . The function  $w$  satisfies the following boundary conditions on  $S_\lambda$ :

$$\frac{\partial^j w}{\partial |x|^j} \Big|_{S_\lambda} = 0, \quad 0 \leq j < s - \frac{1}{2}.$$

Thus (see [15], Chapter 1, Theorem 11.5),  $w$  belongs to  $H_0^s(S_\lambda^i)$ . The extension by zero onto  $S_\lambda^e$  is a bounded operator from  $H_0^s(S_\lambda^i)$  to  $H^s(\mathbb{R}^d)$  (here the condition  $s - (1/2) \notin \mathbb{Z}$  is important — see [15], Chapter 1, Theorem 11.4). Thus, denoting by  $\tilde{w}$  the extension of  $w$  by zero, we get

$$\begin{aligned} \|\tilde{w}\|_{H^s(\mathbb{R}^d)} &\leq C \|w\|_{H^s(S_\lambda^i)} \leq C \|P_i v\|_{H^s(S_\lambda^i)} + C_1 \|P_i v_e\|_{H^s(S_\lambda^i)} \\ &\leq C \|P_i v\|_{H^s(S_\lambda^i)} + C_2 \|v_e\|_{H^s(S_\lambda^i)} \leq C \|P_i v\|_{H^s(S_\lambda^i)} + C_3 \|P_e v\|_{H^s(S_\lambda^e)}, \end{aligned} \quad (8.6)$$

where we have used (8.5) on the last step.

3. Finally, note that  $v = \tilde{w} + v_e$ . This is true on  $S_\lambda^i$  and  $S_\lambda^e$  by the definition of  $v_e$  and  $w$  and thus is also true on  $S_\lambda$  by the embedding theorem (recall that  $s > 1/2$ ). Combining (8.5) and (8.6), we get (8.4). ■

Now without the loss of generality assume that  $\delta + 2\theta \notin \mathbb{Z}$  (otherwise we can slightly increase  $\delta$ ). Thus, by Proposition 8.1, for any  $v \in H^{(1/2)+\delta+2\theta}(\mathbb{R}^d)$ ,

$$\|v\|_{H^{(1/2)+\delta+2\theta}(\mathbb{R}^d)}^2 \leq C (\|P_i v\|_{H^{(1/2)+\delta+2\theta}(S_\lambda^i)}^2 + \|P_e v\|_{H^{(1/2)+\delta+2\theta}(S_\lambda^e)}^2). \quad (8.7)$$

By (8.3) and (8.7), Proposition 7.1 reduces to the consideration of the two VQ,

$$\frac{\|v\|_{H^{(1/2)+\delta+2\theta}(S_\lambda^i)}^2}{\|v\|_{H^{(l/2)+2\theta}(S_\lambda^i)}^2}, \quad v \in H^{(l/2)+2\theta}(S_\lambda^i), \quad (1 - \Delta)^m v(x) = 0, \quad (8.8)$$

$$\frac{\|v\|_{H^{(1/2)+\delta+2\theta}(S_\lambda^e)}^2}{\|v\|_{H^{(l/2)+2\theta}(S_\lambda^e)}^2}, \quad v \in H^{(l/2)+2\theta}(S_\lambda^e), \quad (1 - \Delta)^m v(x) = 0. \quad (8.9)$$

It remains to prove the bound of the type (7.3) for the VQ (8.8) and (8.9).

**3. Reduction of (8.8) to a VQ on  $S_\lambda$ .** For  $s \geq 0$  denote

$$H^{(s)}(S_\lambda) = \bigoplus_{j=0}^{m-1} H^{s-j}(S_\lambda); \quad (8.10)$$

let  $\Gamma_\lambda$  be the restriction operator,

$$\Gamma_\lambda : H^s(S_\lambda^i) \rightarrow H^{(s-\frac{1}{2})}(S_\lambda), \quad v \mapsto v|_{S_\lambda} \oplus \frac{\partial}{\partial|x|}v|_{S_\lambda} \oplus \cdots \oplus \left(\frac{\partial}{\partial|x|}\right)^{m-1}v|_{S_\lambda}. \quad (8.11)$$

For any  $s \in (0, 2m)$  and any  $v$ ,  $(1 - \Delta)^m v = 0$ , one has<sup>1</sup>

$$c\|v\|_{H^s(S_\lambda^i)} \leq \|\Gamma_\lambda v\|_{H^{(s-\frac{1}{2})}(S_\lambda)} \leq C\|v\|_{H^s(S_\lambda^i)}, \quad (8.12)$$

see, e.g., [15], Chapter 2, Theorem 7.4. Thus,

$$\frac{\|v\|_{H^{(1/2)+\delta+2\theta}(S_\lambda^i)}^2}{\|v\|_{H^{(l/2)+2\theta}(S_\lambda^i)}^2} \leq C \frac{\|\Gamma_\lambda v\|_{H^{(\delta+2\theta)}(S_\lambda)}^2}{\|\Gamma_\lambda v\|_{H^{(\frac{l-1}{2}+2\theta)}(S_\lambda)}^2},$$

and so it is sufficient to prove that for the VQ

$$\frac{\|w\|_{H^{(\delta+2\theta)}(S_\lambda)}^2}{\|w\|_{H^{(\frac{l-1}{2}+2\theta)}(S_\lambda)}^2}, \quad w \in H^{(\frac{l-1}{2}+2\theta)}(S_\lambda), \quad (8.13)$$

one has

$$\lim_{s \rightarrow 0} s^\varkappa n(s, (8.13)) < \infty, \quad \varkappa = \frac{d-1}{l-2\delta-1}. \quad (8.14)$$

Consider the VQ's

$$\frac{\|w\|_{H^{\delta+2\theta-j}(S_\lambda)}^2}{\|w\|_{H^{\frac{l-1}{2}+2\theta-j}(S_\lambda)}^2}, \quad w \in H^{\frac{l-1}{2}+2\theta-j}(S_\lambda), \quad (8.15j)$$

for  $j = 0, \dots, m-1$ . One has

$$n(s, (8.13)) \leq \sum_{j=0}^{m-1} n(s, (8.15j)).$$

Applying Proposition 6.1 to each of the VQ's (8.15j), we see that

$$\lim_{s \rightarrow 0} s^\varkappa n(s, (8.8)) < \infty, \quad \varkappa = \frac{d-1}{l-2\delta-1}. \quad (8.16)$$

---

<sup>1</sup>Note that the delicate feature of this estimate is the range of the exponent  $s$ . For  $s \geq 2m$ , the estimate (8.12) is well known and can be found in a number of standard reference texts. However, for  $s \in (0, 2m)$ , the book [15] is the only source of reference we were able to find.

**4. Reduction of (8.9) to a VQ on  $S_\lambda \cup S_{4\lambda}$ .** It remains to prove the estimate of the type (8.16) for the VQ (8.9). We proceed similarly to the proof of (8.16), but the additional technical difficulty to overcome is that the domain  $S_\lambda^e$  is not bounded. Because of this difficulty, we reduce the VQ (8.9) to a VQ on the union of two spheres,  $S_\lambda \cup S_{4\lambda}$  (rather than on the sphere  $S_\lambda$ , as it would seem natural to do). The second sphere  $S_{4\lambda}$  is the price we have to pay for getting rid of the infinity. Our estimates below are by far not optimal, but they are sufficient for our purposes.

Let  $H^{(s)}(S_\lambda \cup S_{4\lambda}) = H^{(s)}(S_\lambda) \oplus H^{(s)}(S_{4\lambda})$  and let us define the restriction operator  $\Gamma : H^s(S_\lambda^e) \rightarrow H^{(s-\frac{1}{2})}(S_\lambda \cup S_{4\lambda})$  by  $\Gamma = \Gamma_\lambda \oplus \Gamma_{4\lambda}$ .

We need an analogue of the estimate (8.12).

**Proposition 8.2.** *For any  $s \in (0, 2m)$  and any  $v$ ,  $(1 - \Delta)^m v = 0$ , one has*

$$c\|v\|_{H^s(S_\lambda^e)} \leq \|\Gamma v\|_{H^{(s-\frac{1}{2})}(S_\lambda \cup S_{4\lambda})} \leq C\|v\|_{H^s(S_\lambda^e)}. \quad (8.17)$$

Using Proposition 8.2, we proceed exactly as for the VQ (8.8) and reduce (8.9) to a ‘direct sum’ of VQ’s on  $S_\lambda \cup S_{4\lambda}$ . After that, using Proposition 6.1, we get

$$\lim_{s \rightarrow 0} s^\varkappa n(s, (8.9)) < \infty, \quad \varkappa = \frac{d-1}{l-2\delta-1},$$

which completes the proof of Proposition 7.1.

**5. Proof of Proposition 8.2:** 1. First note that the same arguments as for (8.12), applied to the domain  $S_\lambda^e \cap S_{4\lambda}^i$ , lead to the estimate

$$c\|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)} \leq \|\Gamma v\|_{H^{(s-\frac{1}{2})}(S_\lambda \cup S_{4\lambda})} \leq C\|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)}$$

for any  $v \in H^s(S_\lambda^e \cap S_{4\lambda}^i)$  such that  $(1 - \Delta)^m v = 0$ . Thus, we only need to prove that for any  $v \in H^s(S_\lambda^e)$ ,  $(1 - \Delta)^m v = 0$ , one has

$$c\|v\|_{H^s(S_\lambda^e)} \leq \|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)} \leq C\|v\|_{H^s(S_\lambda^e)}. \quad (8.18)$$

The second estimate in (8.18) is obviously true. Thus, we only need to prove the first estimate in (8.18).

2. Fix a ‘cutoff function’  $\eta \in C_0^\infty(\mathbb{R}^d)$ ,  $\text{supp } \eta \subset S_{3\lambda}^i$ ,  $\eta(x) = 1$  for  $x \in S_{2\lambda}^i$ . We will prove two estimates:

$$\|\eta v\|_{H^s(S_\lambda^e)} \leq C\|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)}, \quad (8.19)$$

$$\|(1 - \eta)v\|_{H^s(S_\lambda^e)} \leq C\|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)}, \quad (8.20)$$

where  $v \in H^s(S_\lambda^e)$ ,  $(1 - \Delta)^m v = 0$ .

3. Let us prove (8.19). Since  $\eta v$  vanishes near  $S_{4\lambda}$ , one has

$$\|\eta v\|_{H^s(S_\lambda^e)} \leq C\|\eta v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)},$$

cf. [15, Chapter 1, Theorem 11.4]. Next, multiplication by  $\eta$  is a bounded operator in  $H^s(S_\lambda^e \cap S_{4\lambda}^i)$ :

$$\|\eta v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)} \leq C \|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)},$$

so we get (8.19). Note that we have not used the equation  $(1 - \Delta)^m v = 0$ .

4. Let us prove (8.20). Consider the function  $(1 - \eta)v$ , defined on  $S_\lambda^e$ . Denote by  $\tilde{v}$  the extension of  $(1 - \eta)v$  by zero onto  $S_\lambda^i$ . Clearly,

$$\|(1 - \eta)v\|_{H^s(S_\lambda^e)} \leq C \|\tilde{v}\|_{H^s(\mathbb{R}^d)}. \quad (8.21)$$

A direct computation (which uses the equation  $(1 - \Delta)^m v = 0$ ) shows that

$$(1 - \Delta)^m \tilde{v} = \sum_{0 \leq |j| \leq 2m-1} \eta_j D^j v, \quad (8.22)$$

where  $j = (j_1, j_2, \dots, j_d)$  is a multi-index,  $|j| = j_1 + \dots + j_d$ ,  $D^j = \left(\frac{\partial}{\partial x_1}\right)^{j_1} \dots \left(\frac{\partial}{\partial x_d}\right)^{j_d}$ , and  $\eta_j \in C_0^\infty(S_{2\lambda}^e \cap S_{3\lambda}^i)$ . Below we will check the estimate

$$\left\| \sum_{0 \leq |j| \leq 2m-1} \eta_j D^j v \right\|_{H^{s-2m}(\mathbb{R}^d)} \leq C \|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)}. \quad (8.23)$$

Using (8.23) and (8.21), we get

$$\begin{aligned} \|(1 - \eta)v\|_{H^s(S_\lambda^e)} &\leq C \|\tilde{v}\|_{H^s(\mathbb{R}^d)} = C \|(1 - \Delta)^{-m} \sum_{0 \leq |j| \leq 2m-1} \eta_j D^j v\|_{H^s(\mathbb{R}^d)} \\ &\leq C_1 \left\| \sum_{0 \leq |j| \leq 2m-1} \eta_j D^j v \right\|_{H^{s-2m}(\mathbb{R}^d)} \leq C_2 \|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)}, \end{aligned}$$

which proves (8.20).

5. It remains to prove (8.23). To this end, fix another cutoff function  $\beta \in C_0^\infty(\mathbb{R}^d)$ ,  $\text{supp } \beta \subset S_\lambda^e \cap S_{4\lambda}^i$ ,  $\beta(x) = 1$  for  $x \in S_{2\lambda}^e \cap S_{3\lambda}^i$ . One has

$$\sum_{0 \leq |j| \leq 2m-1} \eta_j D^j v = \sum_{0 \leq |j| \leq 2m-1} \eta_j D^j (\beta v).$$

Thus,

$$\begin{aligned} \left\| \sum_{0 \leq |j| \leq 2m-1} \eta_j D^j v \right\|_{H^{s-2m}(\mathbb{R}^d)} &\leq C \sum_{0 \leq |j| \leq 2m-1} \|D^j (\beta v)\|_{H^{s-2m}(\mathbb{R}^d)} \\ &\leq C_1 \sum_{0 \leq |j| \leq 2m-1} \|\beta v\|_{H^{s-2m+|j|}(\mathbb{R}^d)} \leq C_2 \|\beta v\|_{H^s(\mathbb{R}^d)} \\ &\leq C_3 \|\beta v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)} \leq C_4 \|v\|_{H^s(S_\lambda^e \cap S_{4\lambda}^i)}. \quad \blacksquare \end{aligned}$$

## ACKNOWLEDGMENTS

The first author has benefited from discussions with M. Sh. Birman. In particular, he acknowledges with gratitude the fact that he has learnt the key idea of the proof of the estimate (4.11) from M. Sh. Birman and V. A. Sloushch. The authors are grateful to Ya. Kurylev for the discussions concerning Proposition 8.1 and to Yu. Safarov for consultations.

## References

- [1] S. Agmon, *Spectral properties of Schrödinger operators and scattering theory*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **2** (1975), no. 2, 151–218.
- [2] S. Alama, P. A. Deift, R. Hempel, *Eigenvalue branches of the Schrödinger operator  $H - \lambda W$  in a gap of  $\sigma(H)$* , Commun. Math. Phys. **121** (1989), 291–321.
- [3] M. Sh. Birman and V. V. Borzov, *The asymptotic behavior of the discrete spectrum of certain singular differential operators*. (in Russian) Problems of mathematical physics, No. 5: spectral theory, pp. 24–38.
- [4] M. Sh. Birman and M. G. Krein, On the theory of wave operators and scattering operators (in Russian), *Dokl. Akad. Nauk SSSR* **144** (1962), 475–478; English transl. in *Soviet Math. Dokl.* **3** (1962).
- [5] M. Sh. Birman, M. Z. Solomyak, *Compact operators with power asymptotic behaviour of singular numbers*. (in Russian) Zap. Nauchn. Semin. LOMI **126** (1983), 21–30. English translation in: J. Sov. Math. **27** (1984), 2442–2447.
- [6] M. Sh. Birman, M. Z. Solomyak, *Asymptotic behavior of the spectrum of pseudodifferential operators with anisotropically homogeneous symbols*. (in Russian), Vestnik Leningr. Univ. **13**, no. 3 (1977), 13–21. English translation in: Vestnik Leningrad Univ. Math. **10** (1982), 237–247.
- [7] M. Sh. Birman, M. Z. Solomyak, *The principal term of the spectral asymptotics for ‘nonsmooth’ elliptic problems*. (in Russian) Funktsional. Anal. i Prilozhen. **4**, no.4 (1970), 1–13. English translation in: Functional Anal. Appl. **4** (1970), 265–275 (1971).
- [8] M. Sh. Birman, D. R. Yafaev, *The spectral shift function. The papers of M. G. Krein and their further development*. (in Russian) Algebra i Analiz **4**, no. 5 (1992), 1–44. English translation in: St. Petersburg Math. J. **4**, no. 5 (1993), 833–870.

- [9] V.V.Borzov, *Piecewise polynomial approximations of functions of classes  $\overset{\circ}{W}_p^r$  in  $L_q$  with respect to an infinite measure.* (in Russian) Izv. Vys. Uchebn. Zaved. Matematika 1973, **8(135)**, 19–28.
- [10] F. Gesztesy, K. A. Makarov, *The  $\Xi$  operator and its relation to Krein's spectral shift function*, J. Anal. Math. **81** (2000), 139–183.
- [11] I. C. Gohberg, M. G. Krein, *Introduction to the theory of linear nonselfadjoint operators.* Translations of Mathematical Monographs, Vol. 18. AMS, Providence, R.I. 1969.
- [12] M. G. Krein, *On the trace formula in perturbation theory.* (in Russian) Mat. Sb. **33 (75)**, no. 3 (1953), 597–626.
- [13] I. M. Lifshits, *On a problem in perturbation theory.* (in Russian) Uspekhi Mat. Nauk **7**, no. 1 (47) (1952), 171–180.
- [14] A. Horn, *On the singular values of the product of completely continuous operators*, Proc. Nat. Acad. Sci. U.S.A. **36** (1950), 374–375.
- [15] J.-L. Lions, E. Magenes, *Non-homogeneous boundary value problems and applications. Vol. I.* Springer-Verlag, 1972.
- [16] A. B. Pushnitski, *Representation for the spectral shift function for perturbations of a definite sign.* (in Russian) Algebra i Analiz, **9**, no. 6 (1997), 197–213. English translation in: St. Petersburg Math. J., **9**, no. 6 (1998), 1181–1194.
- [17] A. B. Pushnitski, *Spectral shift function of the Schrodinger operator in the large coupling constant limit*, Comm. in PDE **25**, no 3–4 (2000), 703–736.
- [18] A. B. Pushnitski, *The spectral shift function and the invariance principle*, J. Functional Analysis, **183**, no. 2 (2001), 269–320.
- [19] M. Reed, B. Simon, *Methods of modern mathematical physics*, vol. 3, Academic Press, 1979.
- [20] M. Reed, B. Simon, *Methods of modern mathematical physics*, vol. 4, Academic Press, 1978.
- [21] D. Robert, *Semiclassical asymptotics of the spectral shift function*, Adv. in Mathem. Sci. 41; Amer. Math. Soc. Transl. (2) **189** (1999), 187–203.
- [22] O. L. Safronov, *Spectral shift function in the large coupling constant limit*, J. Functional Analysis, **182**, no. 1 (2001), 151–169.

- [23] B. Simon, *Trace Ideals and Their Applications*, Cambridge University Press, Cambridge University Press, Cambridge 1979.
- [24] A. V. Sobolev, *Efficient bounds for the spectral shift function*, Ann. Inst. H. Poincaré Phys. Theor. **58**, no. 1 (1993), 55–83.
- [25] D. R. Yafaev, *Mathematical scattering theory. General theory*, Amer. Math. Soc., Providence, RI, 1992.
- [26] D. R. Yafaev, *Remarks on the scattering theory for a perturbed polyharmonic operator* (in Russian), Mat. Zametki, **15** no. 3 (1974), 445–454.