



Spectral Theory of Orthogonal Polynomials

Barry Simon

IBM Professor of Mathematics and Theoretical Physics
California Institute of Technology
Pasadena, CA, U.S.A.

Lecture 5: Killip–Simon Theorem on $[-2, 2]$

Killip–Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Spectral Theory of Orthogonal Polynomials

Killip–Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

- Lecture 3: Three Kinds of Polynomials Asymptotics, I
- Lecture 4: Three Kinds of Polynomial Asymptotics, II
- Lecture 5: Killip–Simon Theorem on $[-2, 2]$
- Lecture 6: Szegő Asymptotics and Shohat–Nevai for $[-2, 2]$



References

Killip–Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

[OPUC] B. Simon, *Orthogonal Polynomials on the Unit Circle, Part 1: Classical Theory*, AMS Colloquium Series **54.1**, American Mathematical Society, Providence, RI, 2005.

[OPUC2] B. Simon, *Orthogonal Polynomials on the Unit Circle, Part 2: Spectral Theory*, AMS Colloquium Series, **54.2**, American Mathematical Society, Providence, RI, 2005.

[SzThm] B. Simon, *Szegő's Theorem and Its Descendants: Spectral Theory for L^2 Perturbations of Orthogonal Polynomials*, M. B. Porter Lectures, Princeton University Press, Princeton, NJ, 2011.



Killip–Simon Theorem

In 2000, Rowan Killip and I proved the following OPRL analog of Szegő's Theorem.

Theorem. Let $d\mu(x) = f(x) dx + d\mu_s$ with Jacobi parameters $\{a_n, b_n\}_{n=1}^{\infty}$. Then

$$\sum_{n=1}^{\infty} (a_n - 1)^2 + b_n^2 < \infty$$

if and only if

(i) (Blumental–Weyl) $\sigma_{\text{ess}}(J) = \text{ess supp}(d\mu) = [-2, 2]$, i.e., $\text{supp}(d\mu)$ is a set of pure points whose only possible limit points are ± 2 : $E_1^- < E_2^- < \dots < -2$; $2 < \dots < E_2^+ < E_1^+$.

(ii) (Lieb–Thirring) $\sum_{\pm, j} (|E_j^{\pm}| - 2)^{3/2} < \infty$.

(iii) (Quasi-Szegő) $\int (x^2 - 4)^{1/2} \log(f(x)) dx < \infty$.

Killip–Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Killip–Simon Theorem

If J_0 is the Jacobi matrix, $a_n \equiv 1$, $b_n \equiv 0$, the L^2 condition is

$$\mathrm{Tr}((J - J_0)^2) < \infty$$

Weyl's Theorem says $J - J_0$ compact $\Rightarrow \sigma_{\mathrm{ess}}(J) = \sigma_{\mathrm{ess}}(J_0) = [-2, 2]$.

For Schrödinger operators in 1D (and so on half line), Lieb–Thirring proved (initially for $p > 1/2$, $p = 1/2$ is Weidl and then Hundertmark–Lieb–Thomas)

$$\sum_{E_{j,\pm}} |E_j^\pm|^p \leq C_p \int_0^\infty |V(x)|^{p+\frac{1}{2}}$$

Killip–Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Killip–Simon Theorem

Killip–Simon Theorem

P_2 -Sum Rule

Step-by-Step Sum Rules

Step-by-Step \Rightarrow Result

m -Functions

Second Kind Polynomials

Weyl Solution

Meromorphic Herglotz Functions

Case Sum Rules

End of the Story

Hundertmark–Simon (Killip–Simon for $p = 3/2$)

$$\sum (|E_j^\pm| - 2)^p \leq \tilde{C}_p \sum_{n=0}^{\infty} |a_n - 1|^{p+\frac{1}{2}} + |b_n|^{p+\frac{1}{2}}$$

Quasi-Sezgő because power is $+1/2$, not $-1/2$ of Szegő condition.



P_2 -Sum Rule

Define F on $\mathbb{R} \setminus [-2, 2]$ by ($|\beta| > 1$)

$$F(\beta + \beta^{-1}) = \frac{1}{4} [\beta^2 - \beta^{-2} - \log(\beta^4)];$$

$$F(E) = \frac{1}{2} \int_2^{|E|} (y^2 - 4)^{\frac{1}{2}} dy$$

so $F(E) > 0$ and $F(E) = \frac{2}{3} (|E| - 2)^{\frac{3}{2}} + O((|E| - 2)^{\frac{5}{2}})$.

Define $G(a) = a^2 - 1 - \log(a^2)$, so

$G(a) > 0$ on $(0, \infty) \setminus \{1\}$; $G(a) = 2(a - 1)^2 + O((a - 1)^3)$.

$$Q(\mu) = \frac{1}{4\pi} \int_{-2}^2 \log\left(\frac{\sqrt{4-x^2}}{2\pi f(x)}\right) \sqrt{4-x^2} dx$$

$$= -\frac{1}{2} S(\mu_0 | \mu); \mu_0 = (a_n \equiv 1, b_n \equiv 0) \text{ measure}$$

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



P_2 -Sum Rule

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

P_2 -Sum Rule:

$$Q(\mu) + \sum F(E_n^\pm) = \sum_{n=1}^{\infty} \left[\frac{1}{4} b_n^2 + \frac{1}{2} G(a_n) \right]$$

if $\sigma_{\text{ess}}(\mu) = [-2, 2]$.

$$\text{RHS} < \infty \Leftrightarrow \sum_{n=1}^{\infty} b_n^2 + (a_n - 1)^2 < \infty.$$

$$\text{LHS} < \infty \Leftrightarrow \text{Quasi-Szegő} + \sum_{n,\pm} (|E_n^\pm| - 2)^{\frac{3}{2}} < \infty.$$

Thus P_2 -sum rule \Rightarrow KS Theorem.



The Method of Step-by-Step Sum Rules

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

Consider first OPUC. Given μ with Verblunsky coefficients $\{\alpha_n\}_{n=0}^{\infty}$, we define the once stripped measure, μ_1 , by

$$\alpha_j(\mu_1) = \alpha_{j+1}(\mu)$$

i.e., drop α_0 and shift left.

If μ obeys a Szegő condition, so does μ_1 and if $d\mu_1 = f_1 \frac{d\theta}{2\pi} + d\mu_{1,s}$, then

$$1 - |\alpha_0|^2 = \exp\left(\frac{1}{2\pi} \int \log\left(\frac{f(\theta)}{f_1(\theta)}\right) d\theta\right)$$



The Method of Step-by-Step Sum Rules

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

This is a little dicey if f_1 or f vanish on sets of positive Lebesgue measure but otherwise makes sense even if μ doesn't obey a Szegő condition.

There is such a “single step” sum rule in general where $\log \left(\frac{f(\theta)}{f_1(\theta)} \right)$ is replaced by a function $G(\theta)$ equal to that \log if $f(\theta) \neq 0$ and it can be used to prove Szegő's theorem (see [SzThm], Sections 2.6 and 2.7).

Since the proof uses use of entropy, it only replaces variational upper bound so for OPUC not so significant.

Still it leads to a higher-order Szegő theorem for OPUC (see [SzThm], Section 2.8).



The Method of Step-by-Step Sum Rules

We'll eventually prove

Theorem (P_2 Step-by-Step Sum Rule). μ_ℓ has Jacobi parameters $a_j(\mu_\ell) = a_{j+\ell}(\mu)$, $b_j(\mu_\ell) = b_{j+\ell}(\mu)$. Then,

(a) $\sum_{j,\pm} [F(E_j^\pm(\mu)) - F(E_j^\pm(\mu_1))]$ is convergent.

(b) $\exists Q(\mu | \mu_1)$ finite for all μ .

(c) $Q(\mu) < \infty \Leftrightarrow Q(\mu_1) < \infty$ and in that case

$$Q(\mu | \mu_1) = Q(\mu) - Q(\mu_1).$$

$$\frac{1}{4}b_1^2 + G(a_1) = Q(\mu | \mu_1) + \sum_{j,\pm} F(E_j^\pm(\mu)) - F(E_j^\pm(\mu_1))$$

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Step-by-Step Sum Rule \Rightarrow Sum Rule

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

Step 1. P_2 for finite rank perturbations

If $J - J_0$ has rank n , then $\mu_n = \mu_0$ has $Q(\mu_0) = 0 < \infty$.
Thus $Q(\mu) < \infty$. Similarly, the sum of F 's is finite.

By iteration, we get P_2 for $\mu_{n-1}, \mu_{n-2}, \dots, \mu$.



Step-by-Step Sum Rule \Rightarrow Sum Rule

Step 2. Let $J^{(n)}$ have

$$a_\ell^{(n)} = a_\ell, \ell \leq n-1, a_\ell^{(n)} = 1 \text{ if } \ell \geq n,$$

$$b_\ell^{(n)} = b_\ell, \ell \leq n, b_\ell^{(n)} = 0 \text{ if } \ell \geq n+1,$$

Let $\mathcal{E}(\mu) = \sum F$'s.

By Step 1,

$$Q(\mu^{(n)}) + \mathcal{E}(\mu^{(n)}) = \frac{1}{4} \sum_{j=1}^n b_j^2 + \sum_{j=1}^{n-1} G(a_j)$$

Since $Q = -S$, Q is lsc so $Q(\mu) \leq \underline{\lim} Q(\mu^{(n)})$.

For j fixed, $E_j^\pm(\mu^{(n)}) \rightarrow E_j^\pm(\mu)$, so $\sum_{j, \pm \leq m} F(E_j^\pm(\mu)) \leq \underline{\lim} \mathcal{E}(\mu^{(n)})$, so $\mathcal{E}(\mu) \leq \underline{\lim} \mathcal{E}(\mu^{(n)})$.

We have thus proven that $Q(\mu) + \mathcal{E}(\mu) \leq \sum_{j=1}^{\infty} \frac{1}{4} b_j^2 + G(a_j)$.

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Step-by-Step Sum Rule \Rightarrow Sum Rule

Step 3. If $Q(\mu) < \infty$, $\mathcal{E}(\mu) < \infty$, by step-by-step,
 $Q(\mu_n) < \infty$, $\mathcal{E}(\mu_n) < \infty$ and

$$Q(\mu) + \mathcal{E}(\mu) = \sum_{j=1}^{n-1} \left[\frac{1}{4} b_j^2 + G(a_j) \right] + Q(\mu_n) + \mathcal{E}(\mu_n)$$

$$\geq \sum_{j=1}^{n-1} \left[\frac{1}{4} b_j^2 + G(a_j) \right]$$

Taking $n \rightarrow \infty$, $Q(\mu) + \mathcal{E}(\mu) \geq \sum_{j=1}^{\infty} \left[\frac{1}{4} b_j^2 + G(a_j) \right]$

If $Q = \infty$ or $\mathcal{E} = \infty$, this inequality is trivial.

QED!

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



m -Functions

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

One defines $m_\mu(z) = \int \frac{d\mu(x)}{x-z}$ for $z \notin \text{supp}(\mu) = \sigma(J)$.

Of course, m is analytic on $\mathbb{C} \setminus \sigma(J)$ and meromorphic at isolated pure points of μ .

Moreover, since J is multiplication by x in $L^2(\mathbb{R}, d\mu)$, isolated eigenvalues of J are exactly the poles of m_μ .

We'll see soon that the poles of m_{μ_1} , the once-stripped m are precisely the zeros of m_μ .



m -Functions

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

If $(\alpha, \beta) \subset \mathbb{R} \setminus \sigma(J)$, $\frac{dm(y)}{dy} = \int \frac{d\mu(x)}{(x-y)^2} > 0$ so

zeros and poles of m interlace. Since $m \rightarrow 0$ at $\pm\infty$, last “pole or zero” is a pole.

Thus, $E_1^+(\mu) > E_1^+(\mu_1) > E_2^+(\mu) > E_2^+(\mu_1) \dots$

\Rightarrow terms in $F(E(\mu)) - F(E(\mu_1))$ are all positive

and as alternating sum, the sum converges.

Also for Lebesgue a.e. x , $f(x) = \pi^{-1} \lim_{\varepsilon \downarrow 0} \operatorname{Im} m(x + i\varepsilon)$



Second Kind Polynomials

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

**Second Kind
Polynomials**

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

$$q_n(x) = \int \frac{p_n(x) - p_n(y)}{x - y} d\mu(y); \quad q_0 = 0, \quad q_{-1} = -1$$

Since $p_1(x) = a_1^{-1}(x - b_1)$, we have $q_1 = a_1^{-1}$.

Using recursion relation for p 's, see q obeys same relations.

Indeed,

$$q_n(x) = a_1^{-1} p_{n-1}(x; \{a_{\ell+1}, b_{\ell+1}\}_{\ell=0}^{\infty})$$

are “essentially” the p 's for $d\mu_1$.



Weyl Solution

For $z \notin \sigma(J)$, define the Weyl solution

$$g_n(z) \equiv m(z) p_n(z) + q_n(z)$$

which is a solution of difference equation. Thus,

$$\begin{aligned} g_n(z) &= p_n(z) \int \frac{d\mu(x)}{x-z} - p_n(z) \int \frac{d\mu(x)}{x-z} + \int \frac{p_n(x)}{x-z} d\mu(x) \\ &= \langle p_n, (\cdot - z)^{-1} \rangle \end{aligned}$$

Since $(\cdot - z)^{-1} \in L^2(\mathbb{R}, d\mu)$, we see

$$\sum_{n=0}^{\infty} |g_n(z)|^2 < \infty \quad \left(= \frac{\operatorname{Im} m(z)}{\operatorname{Im} z} \text{ if } \operatorname{Im} z \neq 0 \right)$$

If $\inf_n a_n > 0$, the Weyl solution is the unique L^2 solution (up to a constant) by constancy of the Wronskian.

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Weyl Solution

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

$$\text{Clearly } m(z) = -\frac{g_0(z)}{a_0 g_{-1}(z)}$$

since $q_0 = 0$, $p_0 = 1$, $q_{-1} = -1$, $p_{-1} = 0$, $a_0 = 1$.

By uniqueness of L^2 solutions up to a constant

$$g_n(z; d\mu_1) = c(z) g_{n+1}(z; d\mu), n \geq -1.$$

$$\text{Thus, } m(z; d\mu_1) = \frac{-g_1(z)}{a_1 g_0(z)}.$$

In $m(z) = -g_0/a_0 g_{-1}$, we put $a_0 = 1$, but it works for any value of a_0 which is why we put in the a_1 .

Since $a_1 g_1 + (b_1 - z)g_0 + a_0 g_{-1} = 0$, we see that

$$-a_1^2 m_1 + (b_1 - z) - m(z)^{-1} = 0.$$



Weyl Solution

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

Thus, $m(z) = (b_1 - z - a_1^2 m_1(z))^{-1}$, called the coefficient stripping relation.

In particular, poles of m_1 are exactly the zeros of m as we claimed.

Iterating gives Markov continued fraction expansion for m !

In particular taking $z = x + i\varepsilon$, $\varepsilon \downarrow 0$ using $\text{Im}(w^{-1}) = -\text{Im } w / |w|^2$,

$$\varepsilon + a_1^2 \text{Im } m_1 = \text{Im } m / |m|^2 \Rightarrow f/f_1 = |a_1 m|^2$$



Meromorphic Herglotz Functions on \mathbb{D}

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

Let M be meromorphic on \mathbb{D} with $\pm \operatorname{Im} M > 0$ if $\pm \operatorname{Im} z > 0$. Then poles and zeros (i.e., on $(-1, 1)$) interlace. By controlling the ratio of Blaschke products as zeros move, one proves that

Theorem. *If $\{z_j\}_{j=1}^{\infty}, \{p_j\}_{j=1}^{\infty} \subset (-1, 1)$ with $|z_j| \rightarrow 1$ as $j \rightarrow \infty$ and $\sum_{j=1}^{\infty} |z_j - p_j| < \infty$ (automatic if interlaced), then*

$$\prod_{j=1}^N \frac{b_{z_j}(z)}{b_{p_j}(z)} \rightarrow B(z)$$

as meromorphic functions on \mathbb{D} “uniformly” (as functions to Riemann sphere).

B converges in UHP uniformly on compacts, so $|B(e^{i\theta})| = 1$.

As usual $b_{w=0}(z) = z$, $b_{w \neq 0}(z) = -\frac{|w|}{w} \frac{z-w}{1-\bar{w}z}$.



Meromorphic Herglotz Functions on \mathbb{D}

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

**Meromorphic
Herglotz
Functions**

Case Sum Rules

End of the Story

Let B_∞ be Blaschke product of zeros and poles for M , a meromorphic Herglotz function on \mathbb{D} . One proves in UHP, $|\arg B_\infty(z)| \leq 2\pi$ (starting from $\arg B_\infty(x) = 0$ for $B_\infty(x) > 0$ on \mathbb{R}) so $\arg(M/B_\infty)$ is bounded, so by M. Riesz Theorem,

$$\log(M/B_\infty) \in \bigcap_{p < \infty} H^p$$



Meromorphic Herglotz Functions on \mathbb{D}

We get

Theorem. *If M is a meromorphic Herglotz function on \mathbb{D} , B_∞ meromorphic on \mathbb{D} , poles only at poles of M . Then for a.e. θ , $\lim_{r \uparrow 1} M(re^{i\theta}) \equiv M(e^{i\theta})$ exists with*

$$\int \left[\log |M(e^{i\theta})| \right]^p \frac{d\theta}{2\pi} < \infty$$

for all $p \in [1, \infty)$

with

$$f(z) = \sigma B_\infty(z) \exp \left(\int \frac{e^{i\theta} + z}{e^{i\theta} - z} \log |M(e^{i\theta})| \frac{d\theta}{2\pi} \right)$$

where $\sigma = \pm 1$.

$\sigma = \operatorname{sgn}(f(0))$ if $f(0) \neq 0$, $\sigma = 1$ if $f(0) = 0$, $\sigma = -1$ if $f(0) = \infty$.

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Step-by-Step Case Sum Rules

(named after Ken Case) We now apply this to

$$M(z) = m(z + z^{-1})$$

looking at $\log\left(\frac{a_1 M(z)}{z}\right)$. $z = 0$ corresponds to $x = \infty$, so there are Taylor coefficients expressible in terms of continued function expansion. The leading terms are

$$\log \frac{a_1 M(z)}{z} = \log a_1 + b_1 z + \left(\frac{1}{2}b_1^2 + a_1^2 - 1\right)z^2 + O(z^3)$$

Using

$$\log\left(1 - \frac{\beta}{z + z^{-1}}\right) = \sum_{n=1}^{\infty} \frac{2}{n} [T_n(0) - T_n(\frac{1}{2}\beta)] z^n$$

one can obtain “explicit” formulas for the Taylor coefficients.

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story



Step-by-Step Case Sum Rules

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

One also expands the log of Blaschke terms, using

$$\log b_w(z) = \log |w| + \sum_{n=1}^{\infty} \frac{z^n}{n} (w^n - w^{-n})$$

$$\text{and } \frac{e^{i\theta} + z}{e^{i\theta} - z} = 1 + 2 \sum_{n=1}^{\infty} z^n e^{-in\theta}.$$

One gets the C_0 step-by-step rule

$$-\log(a_1) = Z(J | J_1) + \sum_{j, \pm} [\log(|p_j|) - \log(|z_j|)]$$

$$Z(J | J_1) = \frac{1}{4\pi} \int_0^{2\pi} \log \left(\frac{\operatorname{Im} M_1(e^{i\theta})}{\operatorname{Im} M(e^{i\theta})} \right) d\theta$$

if $\operatorname{Im} M(e^{i\theta}) \neq 0$, otherwise it's really $|a_1 M(e^{i\theta})|^2$.



Step-by-Step Case Sum Rules

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

For $n \geq 1$, C_n sum rules,

$$\mathcal{P}_n(a's, b's) = S_n + \mathcal{E}_n$$

\mathcal{P}_n is, in general, complicated but

$$\mathcal{P}_2 = \frac{1}{2} b_1^2 + a_1^2 - 1, \quad \mathcal{P}_1 = b_1$$

$$\mathcal{E}_n = \sum_{j, \pm 1} \frac{z_j^n - p_j^n - (z_j^{-n} - p_j^{-n})}{n}$$

$$S_n = -\frac{1}{2\pi} \int_0^{2\pi} \log \left(\frac{\operatorname{Im} M_1(e^{i\theta})}{\operatorname{Im} M(e^{i\theta})} \right) \cos(n\theta) d\theta$$

where we use $\operatorname{Im} M(e^{i\theta}) = -\operatorname{Im} M(e^{-i\theta})$ so ratio is even to replace $e^{in\theta}$ by $\cos(n\theta)$.



The End of the Story

Killip-Simon
Theorem

P_2 -Sum Rule

Step-by-Step
Sum Rules

Step-by-Step \Rightarrow
Result

m -Functions

Second Kind
Polynomials

Weyl Solution

Meromorphic
Herglotz
Functions

Case Sum Rules

End of the Story

\mathcal{P}_2 is $C_0 + \frac{1}{2}C_2$. A miracle takes place!

$\frac{1}{4\pi} - \frac{1}{4\pi} \cos(2\theta) = \frac{1}{2\pi} \sin^2 \theta$, so the entropies terms combine to

$$Q(J | J_1) = \frac{1}{2\pi} \int_0^{2\pi} \log \left(\frac{\operatorname{Im} M_1}{\operatorname{Im} M} \right) \sin^2 \theta d\theta$$
$$- \log(a_1) + \frac{1}{2} \left(\frac{1}{2} b_1^2 + a_1^2 - 1 \right) = \frac{1}{4} b_1^2 + \frac{1}{2} G(a_1)$$

with $G(a) > 0$ on $(0, \infty) \setminus \{1\}$.

The Blaschke terms also combine to something positive.

Everything works because of the positivity. So far, there is no understanding why they are positive other than as a fortuitous result of calculation!