

This article was downloaded by: [Anthony Coolen]

On: 31 August 2011, At: 13:06

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Philosophical Magazine

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tphm20>

Transfer operator analysis of the parallel dynamics of disordered Ising chains

Anthony C.C. Coolen^{a b} & Koujin Takeda^c

^a Department of Mathematics, King's College London, The Strand, London, WC2R 2LS, UK

^b London Institute for Mathematical Sciences, 22 South Audley St, Mayfair, London, W1K 2NY, UK

^c Department of Computational Intelligence and Systems Science, Tokyo Institute of Technology, 4259 Nagatsuda, Midori-ku, Yokohama 226-8502, Japan

Available online: 31 Aug 2011

To cite this article: Anthony C.C. Coolen & Koujin Takeda (2011): Transfer operator analysis of the parallel dynamics of disordered Ising chains, Philosophical Magazine, DOI:10.1080/14786435.2011.606238

To link to this article: <http://dx.doi.org/10.1080/14786435.2011.606238>



PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan, sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings,

demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Transfer operator analysis of the parallel dynamics of disordered Ising chains[†]

Anthony C.C. Coolen^{ab*} and Koujin Takeda^c

^aDepartment of Mathematics, King's College London, The Strand, London, WC2R 2LS, UK; ^bLondon Institute for Mathematical Sciences, 22 South Audley St, Mayfair, London, W1K 2NY, UK; ^cDepartment of Computational Intelligence and Systems Science, Tokyo Institute of Technology, 4259 Nagatsuda, Midori-ku, Yokohama 226-8502, Japan

(Received 28 March 2011; final version received 14 July 2011)

We study the synchronous stochastic dynamics of the random field and random bond Ising chain. For this model the generating functional analysis method of De Dominicis leads to a formalism with transfer operators, similar to transfer matrices in equilibrium studies, but with dynamical paths of spins and (conjugate) fields as arguments, as opposed to replicated spins. In the thermodynamic limit the macroscopic dynamics is captured by the dominant eigenspace of the transfer operator, leading to a relatively simple and transparent set of equations that are easy to solve numerically. Our results are supported excellently by numerical simulations.

Keywords: disordered Ising chains; parallel dynamics; transfer operator

1. Introduction

In spite of the absence of equilibrium phase transitions at finite temperature in one-dimensional Ising chains, the dynamics of such systems (solved formally several decades ago [1–3]) continue to be of interest in the context of ageing phenomena, see, e.g. [4]. Disordered versions of such chains, with random bonds and/or random fields, generally require new techniques for solution, unless the disorder can be transformed away as for binary bonds. One method for solving disordered chains in equilibrium is based on iteration of partition functions for growing chains, constrained by the state of the last spin [5–9]. More recently such models were also solved by diagonalisation of replicated transfer matrices [10–11]. The situation with the dynamics of disordered Ising chains is less satisfactory. Except for special cases, e.g. [12], our analytical methods are still under development, although it is clear from numerical simulations (e.g. [13,14]) and from the equilibrium solution that the dynamical phenomenology of disordered Ising chains is rich. A renormalisation group approach was advocated in [15]. In this paper, we use the generating functional method of [16] to handle the disorder, and show that this leads to a

*Corresponding author. Email: ton.coolen@kcl.ac.uk

[†]This paper is dedicated to Professor David Sherrington on the occasion of his 70th birthday, to thank him for many years of science and friendship.

transfer operator formalism very similar to that found in equilibrium studies; we use parallel dynamics to keep computations simpler, but results for Glauber dynamics will be similar. While developing our study, another study was published [17], also with parallel dynamics but based on the cavity method, which appears to represent an alternative but mathematically equivalent perspective on some of our equations.

2. Model definitions

We consider N Ising spins $\sigma_i \in \{-1, 1\}$ on a periodic one-dimensional chain. Their dynamics are given by a synchronous stochastic alignment to local fields of the form $h_i(\boldsymbol{\sigma}; t) = J_i\sigma_{i-1} + J_{i+1}\sigma_{i+1} + \theta_i(t)$, with the convention $i + N \equiv i$ for all $i \in \{1, \dots, N\}$, and with $\boldsymbol{\sigma} = (\sigma_1, \dots, \sigma_N)$. Upon defining $p_t(\boldsymbol{\sigma})$ as the probability to find the system at time t in state $\boldsymbol{\sigma}$, this Markovian process can be written as

$$p_{t+1}(\boldsymbol{\sigma}) = \sum_{\boldsymbol{\sigma}'} W_t[\boldsymbol{\sigma}; \boldsymbol{\sigma}'] p_t(\boldsymbol{\sigma}'), \quad W_t[\boldsymbol{\sigma}; \boldsymbol{\sigma}'] = \prod_i \frac{e^{\beta\sigma_i h_i(\boldsymbol{\sigma}'; t)}}{2 \cosh[\beta h_i(\boldsymbol{\sigma}'; t)]}. \quad (1)$$

The parameter $\beta = T^{-1} \geq 0$ measures the noise in the dynamics, which is fully random for $\beta=0$ and fully deterministic for $\beta \rightarrow \infty$. The $\theta_i(t)$ represent external fields of the form $\theta_i(t) = \theta_i + \tilde{\theta}_i(t)$, with random frozen parts θ_i and weak time dependent perturbations $\tilde{\theta}_i(t)$ that serve to define response functions. The bonds J_i and the frozen fields θ_i are regarded as quenched disorder, drawn for each site i independently from a distribution $\tilde{P}(J, \theta)$. We write averages over the process (1) as $\langle \dots \rangle$ and averages over the disorder as $\overline{\dots}$. Upon removing the time dependent parts of the external fields, so that $h_i(\boldsymbol{\sigma}; t) = J_i\sigma_{i-1} + J_{i+1}\sigma_{i+1} + \theta_i$, the process (1) obeys detailed balance, and the equilibrium state will be of the Peretto [18] form

$$p(\boldsymbol{\sigma}) = Z^{-1} e^{-\beta \tilde{H}_\beta(\boldsymbol{\sigma})}, \quad (2)$$

$$\tilde{H}_\beta(\boldsymbol{\sigma}) = - \sum_i \theta_i \sigma_i - \frac{1}{\beta} \sum_i \log 2 \cosh[\beta h_i(\boldsymbol{\sigma})]. \quad (3)$$

The correlation and response functions $C_{ij}(t, t') = \langle \sigma_i(t) \sigma_j(t') \rangle$ and $G_{ij}(t, t') = \partial \langle \sigma_i(t) \rangle / \partial \tilde{\theta}_j(t')$ will be related by the FDT (fluctuation-dissipation theorem) [19]

$$G_{ij}(\tau > 0) = -\beta [C_{ij}(\tau + 1) - C_{ij}(\tau - 1)], \quad G_{ij}(\tau \leq 0) = 0. \quad (4)$$

3. Generating functional analysis

In order to analyse the macroscopic dynamics of the chain we concentrate on the calculation of the disorder averaged generating functional proposed in [16]:

$$\begin{aligned} \overline{Z[\boldsymbol{\psi}]} &= \overline{\left\langle \exp \left[-i \sum_i \sum_{t < t_m} \psi_i(t) \sigma_i(t) \right] \right\rangle} \\ &= \sum_{\boldsymbol{\sigma}(0)} \dots \sum_{\boldsymbol{\sigma}(t_m)} P[\boldsymbol{\sigma}(0), \dots, \boldsymbol{\sigma}(t_m)] \exp \left[-i \sum_i \sum_{t < t_m} \psi_i(t) \sigma_i(t) \right]. \end{aligned} \quad (5)$$

We isolate the local fields at times $t \in \{0, \dots, t_m - 1\}$ in the usual manner via delta functions, using the shorthand $\{\mathbf{d}\mathbf{h} \mathbf{d}\hat{\mathbf{h}}\} = \prod_i \prod_{t < t_m} [dh_i(t)d\hat{h}_i(t)/2\pi]$, which gives

$$\begin{aligned} \overline{Z[\psi]} &= \int \{\mathbf{d}\mathbf{h} \mathbf{d}\hat{\mathbf{h}}\} \sum_{\sigma(0)} \dots \sum_{\sigma(t_m)} p(\sigma(0)) e^{N\mathcal{F}[\{\sigma\}, \{\hat{\mathbf{h}}\}]} \\ &\times \prod_i \prod_{t < t_m} e^{i\hat{h}_i(t)[h_i(t) - \tilde{\theta}_i(t)] - i\psi_i(t)\sigma_i(t) + \beta\sigma_i(t+1)h_i(t) - \log 2 \cosh[\beta h_i(t)]}, \end{aligned} \quad (6)$$

with the disorder dependent exponent

$$\mathcal{F}[\{\sigma\}, \{\hat{\mathbf{h}}\}] = \frac{1}{N} \log \prod_i \int dJ d\theta \tilde{P}(J, \theta) e^{-i \sum_i \{\theta \hat{h}_i(t) + J \hat{h}_i(t) \sigma_{i-1}(t) + \hat{h}_{i-1}(t) \sigma_i(t)\}}. \quad (7)$$

To benefit from the linear nature of the chain we write (6) in terms of the single-site objects $\sigma_i = (\sigma_i(0), \dots, \sigma_i(t_m - 1))$, $\mathbf{h}_i = (h_i(0), \dots, h_i(t_m - 1))$, and $\hat{\mathbf{h}}_i = (\hat{h}_i(0), \dots, \hat{h}_i(t_m - 1))$, with analogous definitions for θ_i and ψ_i . We also introduce the time shift matrix \mathbf{S} , with entries $S_{t,t'} = \delta_{t,t'+1}$, and the vector $\mathbf{u} = (1, \dots, 1)$, so that $\mathbf{x} \cdot \mathbf{S}\mathbf{y} = \sum_t x(t+1)y(t)$ and $\mathbf{u} \cdot \mathbf{x} = \sum_t x(t)$. For factorised and homogeneous initial conditions $p_0(\sigma(0)) = \prod_i p(\sigma_i(0))$ we then obtain

$$\begin{aligned} \overline{Z[\psi]} &= \int \prod_i [d\mathbf{h}_i d\hat{\mathbf{h}}_i] \sum_{\sigma_1 \dots \sigma_N} e^{-i \sum_i [\hat{h}_i \cdot \tilde{\theta}_i + \psi_i \cdot \sigma_i]} \\ &\times \prod_i \langle \sigma_i, \mathbf{h}_i, \hat{\mathbf{h}}_i | M | \sigma_{i-1}, \mathbf{h}_{i-1}, \hat{\mathbf{h}}_{i-1} \rangle, \end{aligned} \quad (8)$$

with operators acting as $(Kf)(\sigma, \mathbf{h}, \hat{\mathbf{h}}) = \sum_{\sigma'} \int d\mathbf{h} d\hat{\mathbf{h}} \langle \sigma, \mathbf{h}, \hat{\mathbf{h}} | M | \sigma', \mathbf{h}', \hat{\mathbf{h}}' \rangle f(\sigma', \mathbf{h}, \mathbf{h}')$, and with a non-symmetric transfer operator M , defined as

$$\begin{aligned} \langle \sigma, \mathbf{h}, \hat{\mathbf{h}} | M | \sigma', \mathbf{h}', \hat{\mathbf{h}}' \rangle &= \frac{p(\sigma(0)) e^{i\hat{\mathbf{h}} \cdot \mathbf{h} + \beta[\sigma \cdot \mathbf{S}\mathbf{h} + \sigma(t_m)h(t_m-1)]}}{\prod_{t < t_m} [4\pi \cosh[\beta h(t)]]} \\ &\times \int dJ d\theta \tilde{P}(J, \theta) e^{-i\theta \mathbf{u} \cdot \hat{\mathbf{h}} - iJ[\hat{\mathbf{h}} \cdot \sigma' + \hat{\mathbf{h}}' \cdot \sigma]}. \end{aligned} \quad (9)$$

Expression (8) is for $N \rightarrow \infty$ dominated by the largest eigenvalue λ_{\max} of (9), provided its spectrum is discrete at λ_{\max} . In an equilibrium replica analysis [10,11] the relevant kernel would have replicated spins as arguments; here the arguments are spin ‘paths’, field ‘paths’ and conjugate field ‘paths’ through time. The fields $\tilde{\theta}_i$ and ψ_i were only introduced for generating perturbations, so we may expand $\overline{Z[\psi]}$ in powers of these fields. To do this efficiently we define

$$\text{Tr}[K] = \int d\mathbf{h} d\hat{\mathbf{h}} \sum_{\sigma} \langle \sigma, \mathbf{h}, \hat{\mathbf{h}} | K | \sigma, \mathbf{h}, \hat{\mathbf{h}} \rangle, \quad (10)$$

$$\langle \sigma, \mathbf{h}, \hat{\mathbf{h}} | S(t) | \sigma', \mathbf{h}', \hat{\mathbf{h}}' \rangle = \sigma(t) \delta(\mathbf{h} - \mathbf{h}') \delta(\hat{\mathbf{h}} - \hat{\mathbf{h}}') \delta_{\sigma, \sigma'}, \quad (11)$$

$$\langle \sigma, \mathbf{h}, \hat{\mathbf{h}} | \hat{H}(t) | \sigma', \mathbf{h}', \hat{\mathbf{h}}' \rangle = \hat{h}(t) \delta(\mathbf{h} - \mathbf{h}') \delta(\hat{\mathbf{h}} - \hat{\mathbf{h}}') \delta_{\sigma, \sigma'}. \quad (12)$$

Since $\overline{Z[\mathbf{0}]} = 1$ for any $\tilde{\theta}$ and $Z[\mathbf{0}] = \text{Tr}[M^N]$ for $\tilde{\theta} = \mathbf{0}$, according to (5) and (8) respectively, we can be sure that $\text{Tr}[M^N] = 1$ and that all terms of order $\tilde{\theta}$ or order $\tilde{\theta}^2$ in our expansion (which can be written in terms of derivatives with respect to $\tilde{\theta}$ of $\overline{Z[\mathbf{0}]}$) must be zero. Thus we may write our expansion in the form

$$\begin{aligned} \overline{Z[\psi]} &= 1 - i \sum_{it} \psi_i(t) \frac{\text{Tr}[S(t)M^N]}{\text{Tr}[M^N]} - \frac{1}{2} \sum_{iit'} \psi_i(t)\psi_{i'}(t') \frac{\text{Tr}[S(t)S(t')M^N]}{\text{Tr}[M^N]} \\ &\quad - \sum_{i < j} \sum_{iit'} \psi_i(t)\psi_j(t') \frac{\text{Tr}[M^{N+i-j}S(t)M^{j-i}S(t')]}{\text{Tr}[M^N]} \\ &\quad - \sum_{i \neq j} \sum_{iit'} \psi_i(t)\tilde{\theta}_j(t') \frac{\text{Tr}[M^{N-|i-j|}S(t)M^{|i-j|}\hat{H}(t')]}{\text{Tr}[M^N]} \\ &\quad + \mathcal{O}(\psi^3, \tilde{\theta}\psi^2, \psi\tilde{\theta}^2). \end{aligned} \quad (13)$$

We may now use the usual relations [16] to express the quantities of interest in the spin chain in terms of derivatives of (5), e.g. $\overline{\langle \sigma_i(t) \rangle} = i \lim_{\psi, \tilde{\theta} \rightarrow 0} \frac{\partial \overline{Z[\psi]}}{\partial \psi_i(t)}$ and $\overline{\langle \sigma_i(t) \sigma_j(t') \rangle} = - \lim_{\psi, \tilde{\theta} \rightarrow 0} \frac{\partial^2 \overline{Z[\psi]}}{\partial \psi_i(t) \partial \psi_j(t')}$, giving

$$m_i(t) = \overline{\langle \sigma_i(t) \rangle} = \text{Tr}[S(t)M^N] / \text{Tr}[M^N], \quad (14)$$

$$C_{ij}(t, t') = \overline{\langle \sigma_i(t) \sigma_j(t') \rangle} = \text{Tr}[M^{N+i-j}S(t)M^{j-i}S(t')] / \text{Tr}[M^N] \quad (i \leq j), \quad (15)$$

$$G_{ij}(t, t') = \lim_{\tilde{\theta} \rightarrow 0} \frac{\partial \overline{\langle \sigma_i(t) \rangle}}{\partial \tilde{\theta}_j(t')} = -i \text{Tr}[M^{N-|i-j|}S(t)M^{|i-j|}\hat{H}(t')] / \text{Tr}[M^N]. \quad (16)$$

Left- and right-eigenvectors with different eigenvalues of (9) are always orthogonal, so we can write (9) in the form $M = \sum_{\lambda} \lambda U(\lambda)$, in which the $U(\lambda)$ are eigenspace projection operators,¹ with $U(\lambda)U(\lambda') = 0$ if $\lambda \neq \lambda'$. The operator $M^\ell = \sum_{\lambda} \lambda^\ell U(\lambda)$ exchanges dynamical information between sites at distance ℓ . Hence $|\lambda| \leq 1$ for all λ , and since $\text{Tr}[M^N] = 1$ for any N , M must have an eigenvalue $\lambda = 1$. Provided the largest eigenvalue is isolated in the spectrum, it follows that $\lim_{N \rightarrow \infty} M^N = \lim_{N \rightarrow \infty} \sum_{\lambda} \lambda^N U(\lambda) = U(1)$, and the above expressions give

$$\lim_{N \rightarrow \infty} m_i(t) = \text{Tr}[S(t)U(1)] / \text{Tr}[U(1)], \quad (17)$$

$$\lim_{N \rightarrow \infty} C_{ij}(t, t') = \text{Tr}[U(1)S(t)M^{j-i}S(t')] / \text{Tr}[U(1)] \quad (i \leq j), \quad (18)$$

$$\lim_{N \rightarrow \infty} G_{ij}(t, t') = -i \text{Tr}[U(1)S(t)M^{|i-j|}\hat{H}(t')] / \text{Tr}[U(1)]. \quad (19)$$

The above quantities are disorder averages of quantities which, by carrying site indices, will not generally be self-averaging. Hence they will not describe the dynamics of an individual realisation of the chain, but averages over many such realisations. In contrast, the following quantities are expected to be self-averaging:

$$m(t) = \lim_{N \rightarrow \infty} N^{-1} \sum_i \langle \sigma_i(t) \rangle = \frac{\text{Tr}[S(t)U(1)]}{\text{Tr}[U(1)]}, \quad (20)$$

$$C(t, t') = \lim_{N \rightarrow \infty} N^{-1} \sum_i \langle \sigma_i(t) \sigma_i(t') \rangle = \frac{\text{Tr}[U(1)S(t)S(t')]}{\text{Tr}[U(1)]}, \quad (21)$$

$$G(t, t') = \lim_{N \rightarrow \infty} N^{-1} \sum_i \lim_{\hat{\theta} \rightarrow 0} \frac{\partial \langle \sigma_i(t) \rangle}{\partial \hat{\theta}_i(t')} = -i \frac{\text{Tr}[U(1)S(t)\hat{H}(t')]}{\text{Tr}[U(1)]}. \quad (22)$$

4. Spectral properties of the transfer operator

From now on we consider only chains with independently distributed bonds and fields, i.e. $\tilde{P}(J, \theta) = \tilde{P}(J)\tilde{P}(\theta)$. This is the natural and technically easier scenario. To study the spectral properties of M it will be helpful to write this operator as

$$\langle \boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}} | M | \boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}' \rangle = (2\pi)^{-l_m} P[\boldsymbol{\sigma} | \mathbf{h}] e^{i\hat{\mathbf{h}} \cdot \mathbf{h}} \int dJ d\theta \tilde{P}(J, \theta) e^{-i\theta \mathbf{u} \cdot \hat{\mathbf{h}} - iJ[\hat{\mathbf{h}} \cdot \boldsymbol{\sigma}' + \hat{\mathbf{h}}' \cdot \boldsymbol{\sigma}]}, \quad (23)$$

with the probability $P[\boldsymbol{\sigma} | \mathbf{h}]$ of a spin exposed to field path \mathbf{h} to follow path $\boldsymbol{\sigma}$:

$$P[\boldsymbol{\sigma} | \mathbf{h}] = p(\sigma(0)) \prod_{t < l_m} \frac{e^{\beta \sigma(t+1)h(t)}}{2 \cosh[\beta h(t)]}. \quad (24)$$

The left- and right-eigenvectors u_R and u_L of (23) are to be solved from, respectively

$$\lambda u_R(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' d\hat{\mathbf{h}}' \langle \boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}} | M | \boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}' \rangle u_R(\boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}'), \quad (25)$$

$$\lambda u_L(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' d\hat{\mathbf{h}}' \langle \boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}' | M | \boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}} \rangle u_L(\boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}'). \quad (26)$$

4.1. Reduction of left- and right-eigenvectors

On the right-eigenvectors u_R we carry out the following transformation:

$$u_R(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) = \int \frac{d\mathbf{x}}{\prod_l (2\pi)} w_R(\boldsymbol{\sigma}, \mathbf{h}, \mathbf{x}) e^{i\hat{\mathbf{h}} \cdot \mathbf{x}} P[\boldsymbol{\sigma} | \mathbf{h}]. \quad (27)$$

Insertion into the eigenvalue equation (25) reveals that $w_R(\boldsymbol{\sigma}, \mathbf{h}, \mathbf{y}) = w_R(\boldsymbol{\sigma}, \mathbf{h} - \mathbf{y})$, and after some trivial manipulations we obtain the simplified eigenvalue problem

$$\lambda w_R(\boldsymbol{\sigma}, \mathbf{h}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' w_R(\boldsymbol{\sigma}', \mathbf{h}') \int dJ d\theta \tilde{P}(J, \theta) \delta[\mathbf{h} - \theta \mathbf{u} - J\boldsymbol{\sigma}'] P[\boldsymbol{\sigma}' | \mathbf{h}' + J\boldsymbol{\sigma}]. \quad (28)$$

Writing out the left-eigenvector equation (26) immediately reveals that $u_L(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) = u_L(\boldsymbol{\sigma}, \hat{\mathbf{h}})$. We now carry out a simple Fourier transformation:

$$u_L(\boldsymbol{\sigma}, \hat{\mathbf{h}}) = \int \frac{d\mathbf{x}}{\prod_l (2\pi)} w_L(\boldsymbol{\sigma}, \mathbf{x}) e^{-i\hat{\mathbf{h}} \cdot \mathbf{x}}. \quad (29)$$

Insertion into the left-eigenvalue problem then gives

$$\lambda w_L(\boldsymbol{\sigma}, \mathbf{h}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' w_L(\boldsymbol{\sigma}', \mathbf{h}') \int dJ d\theta \tilde{P}(J, \theta) \delta[\mathbf{h} - J\boldsymbol{\sigma}'] P[\boldsymbol{\sigma}' | \mathbf{h}' + \theta \mathbf{u} + J\boldsymbol{\sigma}]. \quad (30)$$

The $w_{L,R}(\boldsymbol{\sigma}, \mathbf{h})$ apparently represent distributions of field path contributions, conditioned on spin paths $\boldsymbol{\sigma}$. Given that $\tilde{P}(J, \theta) = \tilde{P}(J)\tilde{P}(\theta)$, they are connected via

$$w_R(\boldsymbol{\sigma}, \mathbf{h}) = \int d\theta \tilde{P}(\theta) w_L(\boldsymbol{\sigma}, \mathbf{h} - \theta \mathbf{u}). \quad (31)$$

To see this we simply define the function $w(\boldsymbol{\sigma}, \mathbf{h}) = \int d\theta \tilde{P}(\theta) w_L(\boldsymbol{\sigma}, \mathbf{h} - \theta \mathbf{u})$ and use (30) to establish that it obeys

$$\begin{aligned} \lambda w(\boldsymbol{\sigma}, \mathbf{h}) &= \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' w_L(\boldsymbol{\sigma}', \mathbf{h}') \int dJ d\theta d\theta' \tilde{P}(J, \theta) \tilde{P}(\theta') \delta[\mathbf{h} - \theta' \mathbf{u} - J\boldsymbol{\sigma}'] P[\boldsymbol{\sigma}' | \mathbf{h}' + \theta \mathbf{u} + J\boldsymbol{\sigma}] \\ &= \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' w(\boldsymbol{\sigma}', \mathbf{h}') \int dJ d\theta \tilde{P}(J, \theta) \delta[\mathbf{h} - \theta \mathbf{u} - J\boldsymbol{\sigma}'] P[\boldsymbol{\sigma}' | \mathbf{h}' + J\boldsymbol{\sigma}]. \end{aligned} \quad (32)$$

Hence $w(\boldsymbol{\sigma}, \mathbf{h})$ obeys (28) and therefore (31) holds. We are now left with only one eigenvalue problem, and upon combining our results we may summarise

$$u_L(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) = \int \frac{d\mathbf{x} e^{-i\hat{\mathbf{h}} \cdot \mathbf{x}}}{\prod_l (2\pi)} \phi(\boldsymbol{\sigma}, \mathbf{x}), \quad (33)$$

$$u_R(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) = P[\boldsymbol{\sigma} | \mathbf{h}] \int \frac{d\mathbf{x} e^{i\hat{\mathbf{h}} \cdot (\mathbf{h} - \mathbf{x})}}{\prod_l (2\pi)} \int d\theta \tilde{P}(\theta) \phi(\boldsymbol{\sigma}, \mathbf{x} - \theta \mathbf{u}), \quad (34)$$

with $\phi(\boldsymbol{\sigma}, \mathbf{x}) \equiv w_L(\boldsymbol{\sigma}, \mathbf{x})$ to be solved from

$$\lambda \phi(\boldsymbol{\sigma}, \mathbf{h}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' \phi(\boldsymbol{\sigma}', \mathbf{h}') \int dJ d\theta \tilde{P}(J, \theta) \delta[\mathbf{h} - J\boldsymbol{\sigma}'] P[\boldsymbol{\sigma}' | \mathbf{h}' + \theta \mathbf{u} + J\boldsymbol{\sigma}]. \quad (35)$$

For $\beta \rightarrow 0$ one easily calculates that $\phi(\boldsymbol{\sigma}, \mathbf{h}) = 2^{-l_m} \sum_{\boldsymbol{\sigma}} \int dJ \tilde{P}(J) \delta(\mathbf{h} - J\boldsymbol{\sigma})$ and that the only possible eigenvalue of (23) is $\lambda = 1$.

4.2. Physical meaning of the $\lambda = 1$ eigenfunctions

The fields experienced at site i can be written as $\mathbf{h}_i = \mathbf{h}_i^R + J_{i+1} \boldsymbol{\sigma}_{i+1} + \theta_i \mathbf{u}$, where $\mathbf{h}_i^R = J_i \boldsymbol{\sigma}_{i-1}$. Apart from the periodicity constraint, all information communicated to site i from spins at sites $j < i$ is channeled via \mathbf{h}_i^R . The conditional likelihood $P_i(\mathbf{h}^R | \boldsymbol{\sigma})$ to observe $\mathbf{h}_i^R = \mathbf{h}^R$ at site i , given we know that $\boldsymbol{\sigma}_i = \boldsymbol{\sigma}$, thus obeys

$$P_i(\mathbf{h}^R | \boldsymbol{\sigma}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' P_{i-1}(\mathbf{h}' | \boldsymbol{\sigma}') P[\boldsymbol{\sigma}' | \mathbf{h}' + \theta_{i-1} \mathbf{u} + J_i \boldsymbol{\sigma}] \delta(\mathbf{h}^R - J_i \boldsymbol{\sigma}'). \quad (36)$$

The spin path $\boldsymbol{\sigma}'$ at site $i-1$ is prescribed in $P_{i-1}(\mathbf{h}' | \boldsymbol{\sigma}')$, so $P_{i-1}(\mathbf{h}' | \boldsymbol{\sigma}')$ no longer depends on θ_{i-1} or J_i . Hence if we average (36) over the disorder we obtain

$$\overline{P_i(\mathbf{h}^R | \boldsymbol{\sigma})} = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' \overline{P_{i-1}(\mathbf{h}' | \boldsymbol{\sigma}') P[\boldsymbol{\sigma}' | \mathbf{h}' + \theta_{i-1} \mathbf{u} + J_i \boldsymbol{\sigma}] \delta(\mathbf{h}^R - J_i \boldsymbol{\sigma}')}. \quad (37)$$

Disorder averaging removes any site dependence of $\overline{P_i(\mathbf{h}|\boldsymbol{\sigma})}$, hence $\overline{P_i(\mathbf{h}|\boldsymbol{\sigma})} = \phi(\mathbf{h}|\boldsymbol{\sigma})$, where the latter is now a true conditional probability distribution, although not corresponding to any specific site, giving the final result

$$\phi(\mathbf{h}|\boldsymbol{\sigma}) = \lim_{N \rightarrow \infty} N^{-1} \sum_i \overline{P_i(\mathbf{h}|\boldsymbol{\sigma})} = \lim_{N \rightarrow \infty} N^{-1} \sum_i \overline{\langle \delta(\mathbf{h} - J_i \boldsymbol{\sigma}_{i-1}) \rangle}_{\boldsymbol{\sigma}=\boldsymbol{\sigma}}, \quad (38)$$

$$\phi(\mathbf{h}|\boldsymbol{\sigma}) = \sum_{\boldsymbol{\sigma}'} \int d\mathbf{h}' \phi(\mathbf{h}'|\boldsymbol{\sigma}') \int dJ d\theta \tilde{P}(J, \theta) \delta(\mathbf{h} - J\boldsymbol{\sigma}') P[\boldsymbol{\sigma}'|\mathbf{h}' + \theta\mathbf{u} + J\boldsymbol{\sigma}]. \quad (39)$$

Equation (39) is identical to (35) for $\lambda = 1$. Expression (38) obeys causality, i.e. $\phi(\mathbf{h}|\boldsymbol{\sigma})$ is independent of $\sigma(t_{\max})$. It is reasonable to assume that for $\lambda = 1$ there is only one solution of (39) that obeys causality, and that non-causal solutions will be ruled out by time boundary conditions. Thus we may for $\lambda = 1$ identify $\phi(\boldsymbol{\sigma}, \mathbf{h}) = \phi(\mathbf{h}|\boldsymbol{\sigma})$. Similar arguments underly the cavity approach in [17], from which (35) is recovered upon substituting the characteristics of the one-dimensional chain.

5. Calculation of observables

To calculate the observables (20)–(21) we need the projection operator $U(1)$. If we make the reasonable assumption that for $\beta > 0$ the $\lambda = 1$ eigenspace is not degenerate, we may use (33), (34), $\text{Tr}[U(1)] = 1$, and $\phi(\boldsymbol{\sigma}, \mathbf{h}) = \phi(\mathbf{h}|\boldsymbol{\sigma})$ to write

$$\begin{aligned} \langle \boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}} | U(1) | \boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}' \rangle &= \gamma^{-1} u_R(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) u_L(\boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}') \\ &= \frac{1}{\gamma} P[\boldsymbol{\sigma}|\mathbf{h}] \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\boldsymbol{\sigma}) \phi(\mathbf{x}'|\boldsymbol{\sigma}') \int d\theta \tilde{P}(\theta) \frac{e^{i\hat{\mathbf{h}} \cdot (\mathbf{h} - \mathbf{x} - \theta\mathbf{u}) - i\hat{\mathbf{h}}' \cdot \mathbf{x}'}}{(2\pi)^{2t_m}}, \end{aligned} \quad (40)$$

with

$$\begin{aligned} \gamma &= \sum_{\boldsymbol{\sigma}} \int d\mathbf{h} d\hat{\mathbf{h}} u_R(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) u_L(\boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}}) \\ &= (2\pi)^{-t_m} \sum_{\boldsymbol{\sigma}} \int d\theta \tilde{P}(\theta) \int d\mathbf{x} d\mathbf{x}' P[\boldsymbol{\sigma}|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'] \phi(\mathbf{x}|\boldsymbol{\sigma}) \phi(\mathbf{x}'|\boldsymbol{\sigma}). \end{aligned} \quad (41)$$

Since $\phi(\mathbf{h}|\boldsymbol{\sigma})$ is independent of $\sigma(t_m)$ and $P[\boldsymbol{\sigma}|\mathbf{h}]$ is independent of $h(t_m)$, we can sum in (41) over $\sigma(t_m)$ and integrate over $h(t_m)$ (in that order), resulting in the same expression for γ but with the replacement $t_m \rightarrow t_m - 1$. Further iteration of this process leads to $\gamma = (2\pi)^{-t_m}$. Hence

$$\begin{aligned} \langle \boldsymbol{\sigma}, \mathbf{h}, \hat{\mathbf{h}} | U(1) | \boldsymbol{\sigma}', \mathbf{h}', \hat{\mathbf{h}}' \rangle &= P[\boldsymbol{\sigma}|\mathbf{h}] \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\boldsymbol{\sigma}) \phi(\mathbf{x}'|\boldsymbol{\sigma}') \\ &\quad \times (2\pi)^{-t_m} \int d\theta \tilde{P}(\theta) e^{i\hat{\mathbf{h}} \cdot (\mathbf{h} - \mathbf{x} - \theta\mathbf{u}) - i\hat{\mathbf{h}}' \cdot \mathbf{x}'}. \end{aligned} \quad (42)$$

We can now write the dynamical observables (20)–(22) in the following physically transparent form (see Appendix A for details):

$$m(t) = \sum_{\boldsymbol{\sigma}} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\boldsymbol{\sigma}) \phi(\mathbf{x}'|\boldsymbol{\sigma}') \int d\theta \tilde{P}(\theta) P[\boldsymbol{\sigma}|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'], \quad (43)$$

$$C(t, t') = \sum_{\sigma} \sigma(t)\sigma(t') \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'], \quad (44)$$

$$G(t, t') = \beta \left\{ C(t, t' + 1) - \sum_{\sigma} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \times \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'] \tanh[\beta(\theta + x(t') + x'(t'))] \right\}. \quad (45)$$

6. Binary bonds and symmetrically distributed random fields

Our equations take a simpler form when the bonds are binary, i.e. for the choice $\tilde{P}(J, \theta) = \tilde{P}(\theta) [\frac{1}{2}(1 + \eta)\delta(J - 1) + \frac{1}{2}(1 - \eta)\delta(J + 1)]$, where $\eta \in [-1, 1]$. Insertion into (39) shows that the dynamic order parameter can now be written as

$$\phi(\mathbf{h}|\sigma) = \sum_{\sigma'} \Phi(\sigma'|\sigma) \delta(\mathbf{h} - \sigma'), \quad (46)$$

$$\Phi(\sigma'|\sigma) = \lim_{N \rightarrow \infty} N^{-1} \sum_i \overline{\langle \delta_{\sigma', J_i \sigma_{i-1}} \rangle}_{\sigma_i = \sigma}, \quad (47)$$

with

$$\begin{aligned} \Phi(\sigma'|\sigma) &= \frac{1}{2}(1 + \eta) \sum_{\sigma''} \Phi(\sigma''|\sigma') \int d\theta \tilde{P}(\theta) P[\sigma'|\sigma'' + \theta\mathbf{u} + \sigma] \\ &\quad + \frac{1}{2}(1 - \eta) \sum_{\sigma''} \Phi(-\sigma''|\sigma') \int d\theta \tilde{P}(\theta) P[-\sigma'|\theta\mathbf{u} - \sigma'' - \sigma]. \end{aligned} \quad (48)$$

If, furthermore, we choose random fields with $\tilde{P}(-\theta) = \tilde{P}(\theta)$ and unbiased initial conditions $p_0(\sigma(0)) = \frac{1}{2}$, then $\int d\theta \tilde{P}(\theta) P[-\sigma'|\theta\mathbf{u} - \sigma' - \sigma''] = \int d\theta \tilde{P}(\theta) \times P[\sigma'|\theta\mathbf{u} + \sigma' + \sigma'']$, and the operator in (39) of which we need eigenfunctions commutes with the spin-flip operator $(F\Phi)(\sigma'|\sigma) = \Phi(-\sigma'|\sigma)$. We then find that $\Phi(-\sigma'|\sigma) = \Phi(\sigma'|\sigma)$, and the relatively simple equation

$$\Phi(\sigma'|\sigma) = \sum_{\sigma''} \Phi(\sigma''|\sigma') \int d\theta \tilde{P}(\theta) P[\sigma'|\theta\mathbf{u} + \sigma + \sigma'']. \quad (49)$$

The formulae for the macroscopic observables can now also be simplified. If we use the following identity, which follows directly from (46), (49),

$$\int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'] = \sum_{\sigma'} \Phi(\sigma|\sigma')\Phi(\sigma'|\sigma), \quad (50)$$

then we find $m(t) = 0$ for all t , and

$$C(t, t') = \sum_{\sigma} \sigma(t)\sigma(t') \sum_{\sigma'} \Phi(\sigma|\sigma')\Phi(\sigma'|\sigma), \quad (51)$$

$$G(t, t') = \beta \left\{ C(t, t' + 1) - \sum_{\sigma} \sigma(t) \sum_{\sigma' \sigma''} \Phi(\sigma' | \sigma) \Phi(\sigma'' | \sigma) \times \int d\theta \tilde{P}(\theta) P[\sigma | \theta \mathbf{u} + \sigma' + \sigma''] \tanh[\beta(\theta + \sigma'(t') + \sigma''(t'))] \right\}. \quad (52)$$

Equations (49), (51), (52) no longer depend on η , since for symmetric field distributions and with $J_i \in \{-1, 1\}$ the observables C and G are invariant under gauge transformations of the type $\sigma_i \rightarrow \tau_i \sigma_i$, so the bonds can be transformed away.

For $\tilde{P}(\theta) = \frac{1}{2} \delta(\theta - \tilde{\theta}) + \frac{1}{2} \delta(\theta + \tilde{\theta})$, with $\tilde{\theta} \geq 0$, we are studying the synchronous dynamics random field Ising chain for which the statics was solved in [9]. If $\tilde{\theta} > 2$ the dynamics close to $T=0$ will be trivial. Each spin freezes into the direction dictated by its external field, and the $T=0$ order parameter reduces to

$$\Phi(\sigma' | \sigma) = \frac{1}{4} \left[\prod_{t=1}^{t_m} \delta_{\sigma'(t), 1} + \prod_{t=1}^{t_m} \delta_{\sigma'(t), -1} \right]. \quad (53)$$

For $\tilde{\theta} < 2$ the dynamics remains non-trivial. Since $P[\sigma | \mathbf{h}]$ and $\Phi(\sigma' | \sigma)$ obey causality, numerical solution of Equations (49), (51), (52) is for binary fields quite manageable.² Examples are shown in Figures 1 and 2, and compared to data from simulations of chains with $N = 10^6$ spins. We observe that $C(t, t') = 0$ for $t - t'$ odd if $\tilde{P}(\theta) = \delta(\theta)$; this is a consequence of a further symmetry of the eigenvalue problem (49), as is shown in Appendix B. For non-zero fields this symmetry is broken, but one still retains a prominent difference between correlations at odd versus even time separations. For those times for which solution of (49), (51), (52) is feasible, i.e. $t, t' \leq 10$, the agreement with simulation data is seen to be perfect.

7. Beyond single-site observables

Let us finally turn to the calculation of observables that involve multiple sites. It is not difficult to transform the eigenvalue problem for M into an equivalent one involving a self-adjoint transfer operator, with the same spectrum. This implies that all eigenvalues λ of M must be real. We note that, unlike the maximum eigenvalue of (35), the non-leading eigenvalues may well depend on the upper time t_m , so we will from now on write $M(t_m)$ and $U(\lambda | t_m)$ instead of M and $U(\lambda)$. Since our formulae for observables cannot depend on which choice is made, provided t_m is equal to or exceeding the largest time argument in the observable, we can combine (17), (18), the expansion $M(t) = \sum_{\lambda} \lambda U(\lambda | t)$, and the property $\text{Tr}[U(1 | t)] = 1$. This allows us to define and work out the two-site correlation function, with $i < j$:

$$\begin{aligned} \tilde{C}_{ij}(t, t) &= \lim_{N \rightarrow \infty} \left\{ \overline{\langle \sigma_i(t) \sigma_j(t) \rangle} - \overline{\langle \sigma_i(t) \rangle \langle \sigma_j(t) \rangle} \right\} \\ &= \text{Tr}[U(1 | t) S(t) M^{j-i}(t) S(t)] - \text{Tr}[S(t) U(1 | t)] \text{Tr}[S(t) U(1 | t)] \\ &= \sum_{\lambda(t) \neq 1} \lambda^{j-i}(t) \text{Tr}[U(1 | t) S(t) U(\lambda | t) S(t)] \\ &\quad + \text{Tr}[U(1 | t) S(t) U(1 | t) S(t)] - \text{Tr}[S(t) U(1 | t)] \text{Tr}[S(t) U(1 | t)]. \end{aligned} \quad (54)$$




Figure 1. Correlations $C(t, t')$ calculated via numerical solution of (49) (for $t, t' \leq 10$, circles) versus correlations measured in numerical simulations with $N = 10^6$ (crosses). Here $T=0.1$ and $\hat{P}(J, \theta) = \frac{1}{2}\delta(\theta)[\eta\delta(J-1) + (1-\eta)\delta(J+1)]$ (which can be mapped onto the non-disordered Ising chain). Left: values of $C(0, t)$, i.e. overlap with the initial state, plotted versus t (which decays as a power law). Right: correlations $C(t, t')$ plotted versus the ratio t/t' , for $t, t' = 1, \dots, 10$. Dots show the values of $C(t, t')$ for larger times $50 \leq t \leq t' \leq 100$, as measured in simulations, showing the typical non-equilibrium behaviour $C(t, t') \sim C(t/t')$ of the ageing regime. Upper branches: even values of $t - t'$; lower branches: odd values of $t - t'$.




Figure 2. Correlations $C(t, t')$ calculated via numerical solution of (49) ($t, t' \leq 10$, circles) versus correlations measured in numerical simulations with $N = 10^6$ (crosses). Here $T=0.1$ and $\hat{P}(J, \theta) = \frac{1}{4}[\delta(\theta - \frac{1}{2}) + \delta(\theta + \frac{1}{2})][\eta\delta(J-1) + (1-\eta)\delta(J+1)]$, i.e. weak random fields. Left: values of $C(0, t)$, i.e. overlap with the initial state, plotted versus t . Right: correlations plotted versus the ratio t/t' , for $t, t' = 1, \dots, 10$. Dots show the values of $C(t, t')$ for larger times $50 \leq t \leq t' \leq 100$, as measured in simulations. Upper branches: even values of $t - t'$; lower branches: odd values of $t - t'$. The system stabilises into a meta-stable state on the time-scales considered, from which it takes significantly more time to escape.

We note that if the correlation length in the chain is finite, then

$$\begin{aligned}
 & \text{Tr}[U(1|t)S(t)U(1|t)S(t)] - \text{Tr}[S(t)U(1|t)]\text{Tr}[S(t)U(1|t)] \\
 &= \lim_{K \rightarrow \infty} \text{Tr}[U(1|t)S(t)M^K(t)S(t)] - \text{Tr}[S(t)U(1|t)]\text{Tr}[S(t)U(1|t)] \\
 &= \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{rs} \left(\overline{\langle \sigma_r(t) \sigma_s(t) \rangle} - \overline{\langle \sigma_r(t) \rangle} \overline{\langle \sigma_s(t) \rangle} \right) \\
 &= \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{rs} \left(\overline{\langle \sigma_r(t) \rangle \langle \sigma_s(t) \rangle} - \overline{\langle \sigma_r(t) \rangle} \overline{\langle \sigma_s(t) \rangle} \right) \\
 &= \lim_{N \rightarrow \infty} \left(\overline{\left\langle N^{-1} \sum_r \sigma_r(t) \right\rangle \left\langle N^{-1} \sum_s \sigma_s(t) \right\rangle} - \overline{\left\langle N^{-1} \sum_r \sigma_r(t) \right\rangle} \overline{\left\langle N^{-1} \sum_s \sigma_s(t) \right\rangle} \right) \\
 &= 0
 \end{aligned} \tag{55}$$

(since the overall magnetisation was assumed to be self-averaging). Hence

$$\tilde{C}_{ij}(t, t) = \sum_{\lambda(t) \neq 1} \lambda^{j-i}(t) \text{Tr}[U(1|t)S(t)U(\lambda|t)S(t)]. \tag{56}$$

If not only $\lambda_{\max} = 1$ but also the second largest eigenvalue $\lambda_2(t)$ is isolated, then the evolving correlation length $\xi(t)$ in the chain can be calculated via

$$\begin{aligned}
 1/\xi(t) &= - \lim_{L \rightarrow \infty} \frac{1}{L} \log \left[N^{-1} \sum_i \tilde{C}_{i, i+L}(t, t) \right] \\
 &= - \lim_{L \rightarrow \infty} \frac{1}{L} \log \sum_{\lambda(t) \neq 1} \lambda^L(t) \text{Tr}[U(1|t)S(t)U(\lambda|t)S(t)] \\
 &= - \log \lambda_2(t).
 \end{aligned} \tag{57}$$

So the present transfer operator picture gives a transparent (although not necessarily trivial) route towards evolving correlation lengths. It is not immediately clear whether and how such formulae could be extracted from the cavity formalism [17].

8. Discussion

We have shown that application of the elegant generating functional analysis method of [16] to disordered Ising chains (with random fields and/or random bonds) leads to a dynamical version of the familiar transfer matrix formalism used in equilibrium studies, with a transfer operator whose arguments are spin paths, field paths and conjugate field paths. Under weak assumptions (e.g. isolated largest eigenvalue in the spectrum of the transfer operator) one can take the thermodynamic limit and find an exact self-consistency equation for a dynamical order parameter, from which disorder-averaged single-site correlation and response functions can be calculated explicitly. The latter equation can also be derived from cavity arguments [17], but without the appealing connection with the equilibrium transfer matrix formalism. As expected, solving the dynamical order parameter equation is still non-trivial. In this paper we have focused on establishing the principles of the method, and we therefore limited ourselves to numerical solution, for short times only. In a future study we

hope to take further steps, and calculate, e.g. equilibrium forms for the order parameter (using the parallel dynamics FDT relation) as well as the form of the solution in the ageing regime. We have also limited ourselves here to investigating those properties than can be extracted from the eigenvectors of the operator that correspond to the largest eigenvalue, which implies calculating single-site objects only. However, in analogy with the usual procedure for equilibrium transfer matrices one can also calculate multi-site quantities (such as evolving correlation lengths) from the second largest eigenvalue of the transfer operator. Whether and how the same could be done within the cavity formalism of [17] is not obvious. Finally, our choice to consider parallel dynamics is not critical for the feasibility of the proposed formalism. In the case of sequential (Glauber) dynamics one will find a very similar structure, but with transfer operators that have continuous time paths rather than discrete time paths as arguments.

Acknowledgements

KT was supported by a Grant-in-Aid (no. 18079006) and Program for Promoting Internationalisation of University Education from MEXT, Japan.

Notes

1. If the spectrum of M has continuous parts, the eigenvalue sum becomes an integral.
2. We interpret (49) as the fixed-point condition of a map $\Phi_{n+1} = (1 - \epsilon)\Phi_n + \epsilon\mathcal{F}\Phi_n$ (where \mathcal{F} denotes the operator in the right-hand side of (49)), and iterate this map until $|\Phi_{n+1} - \Phi_n|$ no longer decreases, upon which ϵ is reduced and the process repeated until a solution of (49) is found. The complexity of solving (49) numerically scales with the dimension of Φ , i.e. as $\sim 2^m$.

References

- [1] R.J. Glauber, *J. Math. Phys.* 4 (1963) p.294.
- [2] B.U. Felderhof, *Rep. Math. Phys.* 1 (1970) p.215.
- [3] B.U. Felderhof, *Rep. Math. Phys.* 2 (1970) p.151.
- [4] C. Godreche and J.M. Luck, *J. Phys. C Condens. Matter* 14 (2002) p.1589.
- [5] P. Rujan, *Physica A* 91 (1978) p.549.
- [6] R. Bruinsma and G. Aeppli, *Phys. Rev. Lett.* 50 (1983) p.1494.
- [7] G. Grinstein and D. Mukamel, *Phys. Rev. B* 27 (1983) p.4503.
- [8] B. Derrida, M. Mendès France and J. Peyrière, *J. Stat. Phys.* 45 (1986) p.439.
- [9] N.S. Skantzos and A.C.C. Coolen, *J. Phys. A Math. Gen.* 33 (2000) p.1841.
- [10] T. Nikolettopoulos, A.C.C. Coolen, I. Pérez-Castillo, N.S. Skantzos, J.P.L Hatchett and B. Wemmenhove, *J. Phys. A Math. Gen.* 37 (2004) p.6455.
- [11] T. Nikolettopoulos and A.C.C. Coolen, *J. Phys. A Math. Gen.* 37 (2004) p.8433.
- [12] G. Forgacs, D. Mukamel and R.A. Pelcovits, *Phys. Rev. B* 30 (1984) p.205.
- [13] H. Rieger, J. Kisker and M. Schreckenberg, *Physica A* 210 (1994) p.324.
- [14] F. Corberi, A. De Candia, E. Lippiello and M. Zannetti, *Phys. Rev. E* 65 (2002) p.046114.
- [15] D.S. Fisher, P. Le Doussal and C. Monthus, *Phys. Rev. E* 64 (2001) p.066107.
- [16] C. De Dominicis, *Phys. Rev. B* 18 (1978) p.4913.
- [17] I. Neri and D. Bollé, *J. Stat. Mech. Theory Exp.* (2009) p.P08009.

[18] P. Peretto, *Biol. Cybern.* 50 (1984) p.51.

[19] A.C.C. Coolen, *Statistical mechanics of recurrent neural networks II: Dynamics*, in *Handbook of Biological Physics*, Vol. 4, F. Moss and S. Gielen, eds., Elsevier Science, Amsterdam, 2001, p.597.

Appendix A. Formulae for magnetisation, correlations, and response function

Here we give details of the derivation of formulae (43)–(45) from (20)–(22). Using $\text{Tr}[U(1)] = 1$ and (42) we can immediately simplify (20), (21) to

$$\bar{m}(t) = \sum_{\sigma} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'], \quad (\text{A1})$$

$$C(t, t') = \sum_{\sigma} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}']. \quad (\text{A2})$$

To evaluate (22) we again use $\text{Tr}[U(1)] = 1$ and (42), followed by the simple identity $\hat{h}(t')e^{i\mathbf{h}\cdot\mathbf{h}} = -i\partial e^{i\mathbf{h}\cdot\mathbf{h}}/\partial h(t')$ and integration by parts over $h(t)$. This gives

$$G(t, t') = \sum_{\sigma} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \int d\mathbf{h} \frac{\partial P[\sigma|\mathbf{h}]}{\partial h(t')} \int d\theta \tilde{P}(\theta) \delta(\mathbf{h} - \mathbf{x} - \mathbf{x}' - \theta\mathbf{u}). \quad (\text{A3})$$

Finally we use

$$\frac{\partial P[\sigma|\mathbf{h}]}{\partial h(t')} = P[\sigma|\mathbf{h}] \frac{\partial \log P[\sigma|\mathbf{h}]}{\partial h(t')} = \beta P[\sigma|\mathbf{h}] [\sigma(t' + 1) - \tanh[\beta h(t')]] \quad (\text{A4})$$

and obtain

$$\begin{aligned} G(t, t') &= \beta \sum_{\sigma} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'] \\ &\quad \times [\sigma(t' + 1) - \tanh[\beta(\theta + x(t') + x'(t'))]] \\ &= \beta \left\{ C(t, t' + 1) - \sum_{\sigma} \sigma(t) \int d\mathbf{x} d\mathbf{x}' \phi(\mathbf{x}|\sigma)\phi(\mathbf{x}'|\sigma) \right. \\ &\quad \left. \times \int d\theta \tilde{P}(\theta) P[\sigma|\theta\mathbf{u} + \mathbf{x} + \mathbf{x}'] \tanh[\beta(\theta + x(t') + x'(t'))] \right\}. \quad (\text{A5}) \end{aligned}$$

Appendix B. Vanishing autocorrelations at odd temporal distance

For $\tilde{P}(\theta) = \delta(\theta)$ Equation (60) simplifies to

$$\Phi(\sigma'|\sigma) = \sum_{\sigma''} \Phi(\sigma''|\sigma') P[\sigma'|\sigma + \sigma'']. \quad (\text{B1})$$

We now define the operator \hat{F} as $(\hat{F}\Phi)(\sigma'|\sigma) = \Phi(\hat{F}^+\sigma'|\hat{F}^-\sigma)$, in which \hat{F}^{\pm} denote commuting spin-flip operators that act on spin states at even (–) or at odd (+) times only:

$$(\hat{F}^+\sigma)(t) = (-1)^t \sigma(t), \quad (\hat{F}^-\sigma)(t) = (-1)^{t+1} \sigma(t). \quad (\text{B2})$$

Clearly $\hat{F}^+ \hat{F}^- = -\mathbf{1}$ and $(\hat{F}^+)^2 = (\hat{F}^-)^2 = \mathbf{1}$. We will prove that Φ is invariant under \hat{F}^+ , and that this causes $C(t, t')$ to be zero whenever $t - t'$ is odd. We note that the function (24) is invariant under \hat{F}^+ , since

$$\begin{aligned} P[\hat{F}^+ \boldsymbol{\sigma} | \hat{F}^- \mathbf{h}] &= p((-1)^0 \sigma(0)) \prod_{t=0}^{t_m-1} \frac{e^{\beta(-1)^{t+1} \sigma(t+1)(-1)^{t+1} h(t)}}{2 \cosh[\beta(-1)^{t+1} h(t)]} \\ &= p(\sigma(0)) \prod_{t=0}^{t_m-1} \frac{e^{\beta \sigma(t+1) h(t)}}{2 \cosh[\beta h(t)]} = P[\boldsymbol{\sigma} | \mathbf{h}]. \end{aligned} \quad (\text{B3})$$

From this it follows upon transforming both sides of (B1) that

$$\begin{aligned} \Phi(\hat{F}^+ \boldsymbol{\sigma}' | \hat{F}^- \boldsymbol{\sigma}) &= \sum_{\boldsymbol{\sigma}''} \Phi(\boldsymbol{\sigma}'' | \hat{F}^+ \boldsymbol{\sigma}') P[\hat{F}^+ \boldsymbol{\sigma}' | \hat{F}^- \boldsymbol{\sigma} + \boldsymbol{\sigma}''] \\ &= \sum_{\boldsymbol{\sigma}''} \Phi(\hat{F}^- \boldsymbol{\sigma}'' | \hat{F}^+ \boldsymbol{\sigma}') P[\boldsymbol{\sigma}' | \boldsymbol{\sigma} + \boldsymbol{\sigma}''] \\ &= \sum_{\boldsymbol{\sigma}''} \Phi((\hat{F}^+)^2 \hat{F}^- \boldsymbol{\sigma}'' | (\hat{F}^-)^2 \hat{F}^+ \boldsymbol{\sigma}') P[\boldsymbol{\sigma}' | \boldsymbol{\sigma} + \boldsymbol{\sigma}''] \\ &= \sum_{\boldsymbol{\sigma}''} \Phi(-\hat{F}^+ \boldsymbol{\sigma}'' | -\hat{F}^- \boldsymbol{\sigma}') P[\boldsymbol{\sigma}' | \boldsymbol{\sigma} + \boldsymbol{\sigma}''] \\ &= \sum_{\boldsymbol{\sigma}''} \Phi(\hat{F}^+ \boldsymbol{\sigma}'' | \hat{F}^- \boldsymbol{\sigma}') P[\boldsymbol{\sigma}' | \boldsymbol{\sigma} + \boldsymbol{\sigma}'']. \end{aligned} \quad (\text{B4})$$

In the last step, we used the previously established symmetry $\Phi(-\boldsymbol{\sigma}' | -\boldsymbol{\sigma}) = \Phi(\boldsymbol{\sigma}' | \boldsymbol{\sigma})$. Hence, if Φ is a solution of (B1) then so is $\hat{F}^+ \Phi$; since the solution of (B1) is assumed to be unique, we deduce that $\hat{F}^+ \Phi = \Phi$. Now

$$\begin{aligned} C(t, t') &= \sum_{\boldsymbol{\sigma}} \sigma(t) \sigma(t') \sum_{\boldsymbol{\sigma}'} \Phi(\boldsymbol{\sigma} | \boldsymbol{\sigma}') \Phi(\boldsymbol{\sigma}' | \boldsymbol{\sigma}) \\ &= \sum_{\boldsymbol{\sigma}} \sigma(t) \sigma(t') \sum_{\boldsymbol{\sigma}'} \Phi(\hat{F}^+ \boldsymbol{\sigma} | \hat{F}^- \boldsymbol{\sigma}') \Phi(\hat{F}^+ \boldsymbol{\sigma}' | \hat{F}^- \boldsymbol{\sigma}) \\ &= \sum_{\boldsymbol{\sigma}} \sigma(t) \sigma(t') \sum_{\boldsymbol{\sigma}'} \Phi(\hat{F}^+ \boldsymbol{\sigma} | \boldsymbol{\sigma}') \Phi(-\boldsymbol{\sigma}' | \hat{F}^- \boldsymbol{\sigma}) \\ &= \sum_{\boldsymbol{\sigma}} (-1)^t \sigma(t) (-1)^{t'} \sigma(t') \sum_{\boldsymbol{\sigma}'} \Phi(\boldsymbol{\sigma} | \boldsymbol{\sigma}') \Phi(\boldsymbol{\sigma}' | \boldsymbol{\sigma}) \\ &= (-1)^{t+t'} C(t, t'). \end{aligned} \quad (\text{B5})$$

Therefore, $C(t, t')$ must vanish if $t - t'$ is odd, as claimed.